SCIENCE LIBRARY

MAY 12 1922

VOL. 49 APRIL, 1922 No. 4

BULLETIN

OF THE

TORREY BOTANICAL CLUB

Editor

ALEXANDER WILLIAM EVANS

Associate Editors

Jean Broadhurst
James Arthur Harris
George T. Hastings
Marshall Avery Howe
Michael Levine
George Elwood Nichols
Arlow Burdette Stout
Norman Taylor

CONTENTS

The phylogeny of the genus Brachiomeeae (Plates 3 and 4)
TRACY E. HAZEN 75
New records and other notes on North Dakota plants.....O. A. STEVENS 93
Three new species of Cuscuta from Mexico.....T. G. YUNCKER 107
INDEX TO AMERICAN BOTANICAL LITERATURE..... 111

PUBLISHED FOR THE CLUB

PRESS OF
THE SCIENTIFIC PRINTING COMPANY
LANCASTER, PA.

THE NEW ERA PRINTING COMPANY

LANCASTER, PA.

Is prepared to execute in first-class and satisfactory manner all kinds of Printing, Electrotyping and Binding. Particular attention given to the work of Schools, Colleges, Universities & Public Institutions.

Books, Periodicals

Technical and Scientific Publications

Monographs, Theses, Catalogues

Announcements, Reports, etc.

All Kinds of Commercial Work

Publishers will find our product ranking with the best in workmanship and material, at satisfactory prices. Correspondence solicited. Estimates furnished.

The New Era Printing Company

American Fern Journal

An illustrated quarterly devoted to the general study of ferns. Subscription, with membership in the AMERICAN FERN SOCIETY, $1.50 per annum; without membership, $1.35; to foreign subscribers, $1.35; life membership in the Society, $25.00. Volume I, six numbers, $2.00; other volumes, $1.00 each. Matter for publication should be sent to R. C. BENEDICT, 332 East 19th Street, Brooklyn, N. Y.; subscriptions and other business communications to E. J. WINSLOW, Auburndale, Mass.

LABORATORY SUPPLIES

OF EVERY DESCRIPTION

MICROSCOPES AND ACCESSORIES

Slides, Coverglasses, Stains, Glassware, Etc.

A well assorted stock always on hand

MEYER CAMERA & INSTRUMENT CO., Inc.

80 Third Avenue NEW YORK

Please mention the "Bulletin" when answering advertisements

Vol. 49 No. 4

BULLETIN

OF THE

TORREY BOTANICAL CLUB

APRIL, 1922

The phylogeny of the genus Brachiomonas

TRACY E. HAZEN

(WITH PLATES 3 AND 4 AND FIVE TEXT FIGURES)

Brachiomonas, one of the oldest and most interesting genera of the *Chlamydomonas* group, has not until now been reported outside of western Europe. The genus was established by Bohlin (**1**) in 1898 with two species, differing in details of form, found in Sweden near Stockholm. The more slender species, *B. gracilis*, has not been reported elsewhere. The type species, *B. submarina*, had been collected about three years earlier by Lagerheim (**6**, p. 7) near Tromsø, Norway, and recorded as a *nomen nudum* which was taken up by Bohlin. A somewhat general distribution of this species on the Norwegian coast has been indicated by Wille (**15**) and Printz (**8**, p. 21). It appears that Dangeard (**3**, p. 74) had observed the same form about ten years earlier than Bohlin, at Luc-sur-Mer, France, but had postponed publishing it until gametes could be found. Chodat (**2**, p. 143, *f. 66*) has recorded the species from Ajaccio, Corsica, showing that it is not exclusively of northern habitat.

West (**12**) in 1908 reported a form collected in brackish marshes at Sheerness, England, near the mouth of the Thames, which he rather hesitatingly referred to *B. submarina*, since it seemed to be somewhat intermediate between Bohlin's two species. In July, 1920, at Cullercoats, near Newcastle-upon-Tyne, and in August at Plymouth, England, I found abundant material in small rock pools, showing forms similar to those described by West, together with specimens agreeing more with Bohlin's type. Earlier in the same season (June 25) I had found on the coast of Norway at Valdersund, near Trondhjem,

[The BULLETIN for March (**49**: 51–74. *pl.* 2) was issued April 12, 1922.]

a few specimens which showed precise gareement with Bohlin's description of *B. submarina*. Again a few days later, at Aalesund, south of Trondhjem, I collected typical material, and here in great abundance. At this station some pools were almost exclusively filled with slender forms corresponding to Bohlin's *B. gracilis*; in other pools a mixture of the two species, with perplexing intermediate forms, appeared. Observing pools containing a natural pure culture, one would scarcely question the distinctness of the two species: when both were found together in the same pool, the variety of forms seen presented a very strong suggestion of hybridization between the two species. New material collected in New York in March and April, 1921, has furnished forms (FIGS. 5–7) almost precisely like the slender cells illustrated by West, together with others showing every gradation to the broadest type of *B. submarina*. These slender forms closely resemble Bohlin's *f. 1b* of *B. gracilis*, but I am now convinced that West was entirely justified in including them in the very variable broader species. We seem not to have in America, and I did not find in England the extreme forms represented by Bohlin's *f. 1a* and *1c* of *B. gracilis*. Whether they are sufficiently distinct to warrant the maintenance of the second species is a question which would be more convincingly settled through studies by pure culture methods.

I first found *Brachiomonas submarina* in March, 1907, in a small rock pool on the shore of Long Island Sound at Twin Island, Pelham Bay, New York. It was found in almost pure culture, coloring the water green from its abundance, and persisted in the motile condition for some weeks when brought to the laboratory. Although the same station was visited many times during the following twelve years, no *Brachiomonas* was found until the middle of November, 1919, when it was again abundant in the same pool, and occurred sparingly in neighboring rock hollows of similar character. I found it repeatedly in the same pools until the first of December, in spite of hard frosts and ice formation during these two weeks. The habitat appears to be very like that of Bohlin's type station, small hollows so high above ordinary tide limits as to seem to be filled only with rain water, which are nevertheless sometimes dashed by waves so as to become brackish, as evidenced by taste and by the salt incrustation left on the rock margin upon evaporation. About the time of the collection in 1919, an unusually high tide was reported, and it is possible that this may have washed the

Brachiomonas into the pool from the open sea. Such an occurrence might account for the sporadic appearance of the species at this station, though it is much more probable that it is usually present in a resting state and that conditions were not favorable for the production of zoospores at the time of other visits to the station. A strange point is that, although careful examination of the specimens collected in the spring of 1907 shows them all to be like Bohlin's type, as were also the few specimens found there in September, 1920, and abundant material from March to June and in September, 1921, only a few such individuals were found in the fall of 1919 (FIGS. 1, 2), while most of the material showed a distinct difference in form (FIGS. 8–16). This can hardly be interpreted as a mere seasonal variation. New material collected the last of February of the present season, from pools still coated with ice, consisted very largely of the new obtuse form, but after a week or more in the laboratory this form had mostly been replaced by individuals corresponding to Bohlin's type, present only in small numbers at the time of collection. Whether this is a case of the new type actually giving rise to the standard type through reproduction, or merely an example of the regular succession of growth of different organisms familiar in collections of protozoa, is a matter calling for more extended observation. For the determination of the permanent status of the new form, pure culture methods are highly desirable, but thus far I have been unable to grow *Brachiomonas* on seawater agar, which succeeds admirably with other chlamydomonads of similar habitat. For the present it will be convenient to discuss this divergent plant under the name

BRACHIOMONAS SUBMARINA Bohlin, forma **obtusa** f. nov.

This new form is entirely like Bohlin's type in its genera shape; that is, it possesses a long posterior extension or horn and four shorter median arms directed backwards, often with depressions between them; the anterior portion might be roughly likened to the shape of an umbrella with four ribs. There is often a wedge-shaped beak through which the cilia emerge, though sometimes the apex appears merely rounded. The protoplast may fill the arms and posterior horn entirely (FIG. 8) or partially (FIG. 12) or not at all. The massive chromatophore may be equally variable; that is the protoplasm in the horn and in the arms may be all green, as also in the typical form (FIG. 2), or it may be colorless, sometimes appearing sharply

delimited from the central green mass (FIG. 8), while in other cases the green and colorless regions merge into one another almost imperceptibly (FIG. 1). The chromatophore is here certainly less definitely or constantly organized than in most species of *Chlamydomonas*; in fact it would perhaps be more accurate not to speak of a definite chromatophore but merely of cytoplasm with chlorophyll more or less diffused through it. This variability in the extent of the distribution of the chlorophyll in the protoplasm is very similar to the condition in *Hydrodictyon*, to which Harper (**5**, p. 179) has called attention. West (**12**) states that in his specimens of *B. submarina* the chromatophore never extended into the arms. This was undoubtedly due to the fact that he never saw his material when freshly collected, but only after it had been sent from near the mouth of the Thames to Birmingham. I find that very generally in freshly collected specimens the chromatophore fills the arms and posterior horn, but often retracts from them soon after being subjected to the less favorable light and aeration conditions of captivity.

Imbedded in the chromatophore in a subparietal position, regularly on the side next to the stigma, is a single relatively large pyrenoid; in vigorous young specimens this is generally situated in front of the middle of the cell; often in older individuals it becomes posterior in position. The conspicuous red eye-spot is linear in form; or, sometimes at least, it may be very narrowly wedge-shaped, pointed in front and broader behind; it lies in the depression midway between two arms. Most frequently when the cell is at rest the pyrenoid and stigma lie in a median ventral position (FIG. 6). On the other side of the cell from the pyrenoid the nucleus may sometimes be detected without staining, lying in a colorless mass of cytoplasm, best seen in polar view (FIG. 13). When the pyrenoid lies in its characteristic anterior position the nucleus is likely to be somewhat posterior (FIGS. 15, 17); if the pyrenoid becomes posterior the nucleus is then more often median in position. In older stages vacuoles often develop in the chromatophore (FIG. 4) as described by West (**12**).

In hanging drop cultures, or in jars kept for some time in the laboratory, many individuals may be found which show little or no trace of the four lateral arms (FIGS. 14, 15). Bohlin has shown such a form in one dividing individual. The cell wall is of so firm a character that it is difficult to imagine how the arms

could be modified or suppressed after they are once formed; nevertheless in practically all examples of division seen, I have found the arms well developed before the escape of the daughter cells (FIGS. 20, 21) so that such cells can hardly be looked upon as a reversion to the simpler ancestral form.

The main difference between this new form and Bohlin's type lies in the fact that the arms are here always rounded and obtuse, while in the type form they are sharp pointed and often more slender. The characters of the form may be summarized as follows:

BRACHIOMONAS SUBMARINA Bohlin, forma **obtusa** f. nov. *B.* cellulis vegetativis figura structuraque formae typicae similibus, sed brachiis cornuque posteriore obtusis; cytoplasmate et chromatophoro alias in extrema ipsa extendentibus, alias ex eis plus minusve retractis.

Longit. cellularum 15–32μ (saepius 20–25μ); lat. cell, 15–22μ (saepius ca. 18μ).

Hab. in aquis subsalsis lacunarum saxearum. Twin Island, Pelham Bay, New York; Nov., 1919, Feb.–March, 1922.

Chodat's figures (**2**, *f. 66*) indicate that the form collected in Corsica may be identical with the one here described, though it is possible that the difference between his sketches and those of Bohlin may be due to the personal equation in drawing.

As indicated above, I have found this obtuse form only at one station. Bohlin's type, as represented by my FIG. 1–3, I have also collected early in September during each of the past two years at Bass Point, Nahant, Massachusetts. It should be looked for on the coast of Maine and New Brunswick as well, at any points where rocky ledges occur.

In older cultures numerous cells were found which showed the entire protoplast contracted into a globose mass (FIG. 16). These showed no further development in this material; but in fresh hanging drop cultures of *B. submarina* followed for a much shorter time in Norway, such cells presently became quiescent and developed a new spherical wall closely investing the protoplast. Soon these aplanospores showed their chromatophore turning from green to a tawny tint, and finally the original wall of the zoospores gradually disappeared.

Practically all genera allied to *Chlamydomonas* possess two contractile vacuoles pulsating more or less in alternation, situated near the base of the cilia. In *Brachiomonas* these appear to be entirely absent. One of West's figures (**12**, *pl. 20, f. 11*; **14**, *f. 101 c*), it is true, suggested such vacuoles, but his

beautiful original sketches in color, preserved with his collection of drawings in the South Kensington Museum, show that these spots are brown colored granules suspended in the fluid of a large permanent vacuole. The absence of contractile vacuoles I have been inclined to regard as another mark of the highly specialized character of the genus, since they are regularly to be expected in primitive members of this group and its ancestors. It is possible, however, that this peculiarity is merely an incident due to the special habitat; for Bütschli has remarked the general absence of contractile vacuoles in marine Ciliates and Rhizopods, and D'Arcy Thompson (**11**, pp. 165 and 264) regards this absence in such cases as "no more than a physical consequence of the different conditions of existence in fresh water and in salt." The marine species, *Chlamydomonas brachyura* West, besides exhibiting other points of resemblance to *Brachiomonas*, appears also to have no contractile vacuoles; this species was discovered in association with a new Carteriaceous genus *Platymonas*, in which West (**13**) found two small contractile vacuoles, which, however, "could only be observed with difficulty." In two brackish water species of *Platymonas* which I have investigated with considerable care, I am unable to satisfy myself that I ever see contractile vacuoles. Lewis has just published figures of one of these species, *P. subcordiformis* (Wille) Hazen,* which likewise show no contractile vacuoles. Furthermore, two new brackish water forms of the primitive family Polyblepharidaceae, one a minute species of *Pyramimonas* and the other a new genus most closely related to *Polyblepharides*, which have turned up in my latest collections of *Brachiomonas*, made while this paper was in the hands of the printer, very certainly possess no contractile vacuoles. Over against these several records of their absence, however, attention may be called to their presence (though here very small) in *Chlamydomonas caudata* Wille, discussed below, which must be considered as regularly a marine organism.

Asexual reproduction is accomplished by internal division into four or eight (very rarely perhaps two) daughter cells. The division very generally begins while the cell is swimming actively, and often the typical cell shape is acquired by the daughter cells or zoospores and they move with their own cilia inside the mother cell-wall while it continues in active movement

* Notes from the Woods Hole Laboratory,—1921. *Platymonas subcordiformis* (Wille) Hazen. Rhodora **23**: 249–251. *pl. 133, f. 1–19.* Mr 1922.

by its own cilia. Bohlin states that the first division is longitudinal, and certainly this is regularly the case. Chodat presents a figure (**2**, *f. 66*), and I have seen two or three such cases, where the first division appears to be transverse, but it is quite possible that such an appearance is due to an unusual shifting of the position of the cell contents after a proper longitudinal division. The subsequent divisions have not been described or figured by previous observers, and are ordinarily difficult to follow because of the continual active movement of the mother cell. I was so fortunate as to find one very favorable case in *B. submarina* f. *obtusa*, and later many similar ones in the type form, which I believe to be entirely representative. This individual became quiescent after the first division was completed (FIGS. *22*, *23*); it presented its anterior pole upward, so that it was possible to see easily that the second plane of cleavage is also longitudinal and perpendicular to the first. Immediately after the second cleavage a slight stretching and shifting of the daughter portions took place, so that one of them appeared somewhat underneath two others (FIG. 24). The third cleavage appears to cut each of the four daughter portions across its longer diameter, that is the plane seems to be essentially transverse to the previous divisions and to the axis of the mother cell. The pyrenoid disappears before the first cleavage occurs, and the stigma fades out during this process. A new and rather small pyrenoid appears in each daughter portion after the second cleavage, and probably another pyrenoid arises *de novo* in four of the eight daughter cells. About an hour after the completion of the third cleavage the eight daughter cells had acquired the typical form of the arms, and also the eyespot and cilia (FIG. 27); after about another hour they escaped from the mother cell-wall. In certain cases it is clear that the cilia of the mother cell remain connected by a strand of protoplasm to one of the daughter portions during division (FIG. 19) or even until the eight zoospores acquire their final form. Whether this is regularly the case appears doubtful. In material identified as *Brachiomonas gracilis* Bohlin, at Aalesund, Norway, I observed numerous cases of division which followed the course outlined above.

In the case of the production of only four daughter cells it is clear that the cytoplasmic polarity of the mother cell would be directly transferred to each of the new zoospores. But when division proceeds further it is less clear how the

polarity may be transmitted: for, as stated above, the third plane appears to cut the four daughter portions transversely in such a manner that half of the resulting eight would retain the original anterior cytoplasm, while the four posterior zoospores would seem to be under the necessity of regenerating (so to speak) their anterior pole. It is possible that in the slight shifting of the daughter portions following so quickly upon the first two divisions there is actually more rotation than is apparent so that the third division may effect a properly qualitative bipartition, but that appears doubtful from the cases observed. The whole problem of polarity in chlamydomonads is one of great interest demanding a degree of care in observation which hardly any species up to the present has received.

Brachiomonas simplex sp. nov.

This species was first discovered at Aalesund, Norway, in the last days of June 1920, and was also collected three weeks later at Cullercoats, near Newcastle-upon-Tyne, and in August at Plymouth, England. It was found in small rock pools of brackish water, sometimes in association with one of the other species of *Brachiomonas*, then again in practically a natural pure culture.

At first sight the organism looked like a *Chlamydomonas*, many individuals (FIGS. 31, 39) showing somewhat the general oval, posteriorly pointed form characteristic of *C. caudata* Wille (FIGS. 46–50), which occurred near by, though never in the same pools. But presently it was noticed that even the cells most nearly approaching *C. caudata* in form were nevertheless more like *Brachiomonas submarina* in internal organization, showing a similar linear stigma and large lateral and anterior pyrenoid; furthermore many individuals show small bumps or protuberances slightly back of the middle region, immediately suggesting the arms of *Brachiomonas*, though never really attaining that character (FIGS. 28, 29). The posterior horn, often rather elongated, shows a characteristic curve not seen in *B. submarina*. Generally the protoplast was conspicuously furnished with vacuoles, but whether this is a constant character is somewhat doubtful, inasmuch as *C. caudata* in similar pools was often similarly vacuolate at this time. The English material, moreover, was much less vacuolate. There is great variability in the development of the lateral protuberances; if formed at all they appear before the escape of the zoospores

from the mother cell, and the form then attained persists through the life of the individual. In newly formed zoospores, usually, but not always, the protoplast and chromatophore extend into the posterior horn so as to fill it completely. In older individuals the protoplast gradually retracts so that a considerable portion of the horn, or finally all of it is empty. The pyrenoid and red eye-spot regularly lie on the same side of the cell, and the nucleus on the opposite side; as in *B. submarina*, this is most clearly shown in cells momentarily resting with one of the poles directed upward (FIG. 32).

As in the other species, asexual reproduction is accomplished by division of the protoplast into four or eight daughter cells which develop the characteristic form while the mother cell still retains its cilia in active movement (FIGS. 35, 36).

In sexual reproduction, gametes may be formed to the number of thirty-two in one mother cell, which retains its motile condition until they are completely developed (FIG. 40) just as in the case of asexual reproduction. The gametes are similar in form to the asexual cells, but much smaller, lacking a cell wall, and furnished with relatively longer cilia: they also seemed to be more apt to show angularity in the middle region than the asexual cells. Usually there was a slight differentiation between the pair, a smaller and more fusiform gamete conjugating with one slightly larger and more angular (FIGS. 41, 42). Meeting by the ciliated anterior ends, the fusion is lateral, resulting in a quadriciliate zygote of ovoid-pyriform shape which swims actively, and may for some time show the two nuclei, two pyrenoids, and two linear eye-spots (FIG. 43); when momentarily at rest the four long cilia trail backwards. Finally the nuclei fuse, the cilia disappear (FIG. 44) the form becomes spherical, and the thick wall of the resting zygospore is developed (FIG. 45). After some days the cell contents assume a reddish orange color similar to that of the aplanospore of this species and of *B. submarina*.

Brachiomonas simplex sp. nov. *B.* cellulis vegetativis majoribus, plerumque oviformibus quidem sed ad posteriorem partem in cornu curvatum productis, brachiis quae ceterae hujus generis species habent aut omnino suppressis aut evolutis tantummodo in 2–4 gibbos exiguuos subangulares vel rotundatos; ad polum autem anteriorem globosis vel papilla conica parvula exornatis: chromatophoro cellulam prope complente aut non longe in cornu posterius se extendente, pyrenoidem lateralem modo in medio modo a fronte habente, saepe vacuola majuscula

plura praebente: puncto rubro (stigmate) lineari prope pyrenoidem sito: nucleo ad latus oppositum cellulae intra partem chromatophori excavatam complexo: ciliis binis ca. ¼ brevioribus quam cellula instructis.

Propagatio fit per cellulae vegetativae impigre natantis divisionem in 4 vel 8 zoosporas, quae tegumenti scissura erumpunt. Praeterea aplanosporae, primo virides, postea paulatim ad colorem fulvum transmutatae, per contractionem protoplasti in massam globosam ad cellulae partem anteriorem, ut in *B. submarina*, efficiuntur.

Generatio fit per copulationem gametarum inter se subaequalium et nudarum, quae cellulis vegetativis forma quidem similes sunt, sed perparvae et ciliis quam corpus ipsum multo longioribus praeditae, quaeque per divisionem cellulae vegetativae protoplasti etiam in 32 individua parturiuntur. Zygosporae globose, tegumento levi vestitae, diametro ca. 12μ.

Longit. cellularum veg. 30–48μ, lat. 18–24μ. Longit. gametarum 13–15μ, lat. 6–8μ; longit. ciliorum ca. 18μ.

Hab. in aquis lacunarum saxearum subsalsis. Aalesund, Norway, 28 June, 1920; Cullercoats, Northumberland, England, 19 July, 1920; Plymouth, England, 11 August, 1920.

Because of the fact that unarmed individuals appear in older cultures of *B. submarina*, one might be tempted to inquire whether *B. simplex* is merely a form of that species. Nevertheless in the natural pure cultures seen both at Aalesund and at Plymouth, *B. simplex* maintained its characteristic simplicity of form from the earliest development of the daughter cells to maturity, and even when it was found intermingled with one of the other species it showed a distinctive character, particularly in the curving of the posterior horn, which rendered it always recognizable. Again, the variability of form exhibited by this new species might readily suggest for it a hybrid origin. We know little of hybridization among algae, but a very interesting preliminary report of its occurrence in *Chlamydomonas* has recently been published by Pascher (**7**). It may be remarked that Pascher's heterozygous cells showed great diversity in internal organization, while here in *B. simplex* only variability of external form is found. From a careful consideration of all the conditions of its habitat and its behavior, I am convinced that *B. simplex* is to be regarded as a distinct species which represents the transition from *Chlamydomonas* to the more characteristic species of *Brachiomonas*.

There is some resemblance between this species and *Chlamydomonas brachyura* West (**13**) a form which developed in a tank of seawater sent from Plymouth to Birmingham in 1915, and for

which I searched vainly at Plymouth in 1920. Nevertheless since *C. brachyura* lacks a red eye-spot and is pointed instead of rounded at the anterior end, we may perhaps find more suggestion of the origin of *Brachiomonas* in the species mentioned at the beginning of this section, namely,

CHLAMYDOMONAS CAUDATA Wille

This very interesting and apparently rare species was found June 28, 1920, at Aalesund, Norway, in small rock pools of the same character as indicated in the original account in 1903 (**15**, pp. 115–118, 135, *pl. 3, f. 4–11*) and my determination of the species was later confirmed by Professor Wille, from living material which I carried to him in Christiania. In one pool was a practically pure culture, giving an intense green color to the brackish water; in another pool close by was a considerable admixture of *C. subcaudata* Wille, which was certainly the most abundant species of the region, now as in 1902. Why *C. caudata* should occur in only three or four out of many similar pools is an interesting question. It was also surprising that it could not be found on the English coast, where the conditions appeared to be much the same, and where I did find several of the other brackish-water species which were associated with *C. caudata* at Aalesund. This species appears not to have been found elsewhere than at this original station.*

The elongated oval posteriorly pointed form of the zoospores is very notable in comparison with the rounded form of the commonly known species of *Chlamydomonas*. The strong resemblance to the more reduced forms of *Brachiomonas simplex* is equally striking; so much so, in fact, as intimated above, that upon superficial observation the two species might be mistaken for closely related members of the same genus. Ordinarily in active individuals, especially in newly formed zoospores, the protoplast conforms rather closely to the shape of the cell-wall, being merely rounded at the point of insertion of

* Playfair, to be sure, has reported [Proc. Linn. Soc. New South Wales **43**: 515. 1918] a form found apparently in fresh water of which he writes: "The only tailed species of *Chlamydomonas*; it is impossible to make any mistake in the identification." Since, however, Playfair states that his specimens lacked pyrenoid and stigma, and his figure shows a distinctly different morphology of the anterior portion of the cell, including a different position of the nuoleus, there can be no justification for trying to identify his form with a species so well defined as *C. caudata*, at least until cultural evidence is brought forward to support the attempt.

the cilia, and tapering to an acute point at the posterior end FIG. 46). Older individuals, kept for two weeks or longer in hanging drops, show the cell-wall much thickened, or the protoplast contracted and rounded posteriorly and produced into more or less of a beak at the anterior end (FIGS. 47, 48). The broad truncate anterior papilla of the cell-wall (Hautwarze) impresses me as being rather more prominent than shown in



FIGS. 46–50. CHLAMYDOMONAS CAUDATA Wille

46. Young cell, "dorsal view," showing typica lposition of nucleus, stigma, and pyrenoid. 47. Older cell with thickened wall. 48. Similar cell in "lateral view." 49. Typical arrangement of daughter cells. 50. Unusual arrangement of daughter cells. All, × 720, approximately.

Wille's (15) figures, and the red eye-spot, first described as 'langgestreckter' and again as 'oval oder stabförmig,' was in my mature specimens always a strongly oval disc, sometimes with an anterior pointed end, showing its thinness in the fact that the edge view was narrowly linear. These differences may be readily accounted for by differences of magnification during observation. The longitudinal streaking of the chromatophore noted in the original description, and even introduced in Wille's key to the species, was never distinguishable in vigorous material of either this species or *C. subcaudata* in 1920; only certain individuals kept for ten days or more in

hanging drop cultures showed almost a lattice-like appearance of the chromatophore. Although the chromatophore is hollowed out about as far as the middle of the cell, or sometimes even more deeply, nevertheless the position of the nucleus is notably less central than might be expected, for it lies approximately in the anterior third of the cell.

A point of considerable interest is the position of the stigma and contractile vacuoles. Regularly the eye-spot is so placed that when the cilia are stretched out in a plane parallel to the stage of the microscope the stigma lies on the apparent right side of the cell and in a slightly upper focus, or with the cell revolved 180° the stigma lies on the apparent left side and at a slightly lower focus; in such a position only one of the small contractile vacuoles is visible. If the cell is revolved only 90° from the first position, so that one of the cilia lies nearer the observer and the other at a somewhat lower focus, the stigma shows its disc-like shape in an upper central focus (FIG. 48) and the two contractile vacuoles may be seen side by side at the same time, though one is always smaller than the other, on account of the alternation in their pulsations. This position of the stigma and cilia in one plane and the contractile vacuoles in another plane perpendicular to it indicates a dorsi-ventral character of the cell which appears to me to be the rule in many species of *Chlamydomonas*, though it seems to have received little or no attention from students of the group.*

The dorsiventral differentiation in *Brachiomonas*, as mentioned above and indicated in FIGS. 6, 7, 13, 15, and 32, is slightly different; there the cilia normally stretch out in a plane perpendicular to that which passes through the stigma, pyrenoid, and nucleus, and most commonly the zoospore comes to rest with the stigma lying exactly in the middle on the ventral side, or (less often) on the dorsal side, toward the observer. In many of the figures which seem to indicate a different relative position of the eye-spot and cilia, the latter were actually lying in a plane oblique to that of the microscope stage, which could not easily be shown in the drawings.

* There are, it is true, many published figures of *Chlamydomonas* which show the two contractile vacuoles side by side apparently in the same plane with the cilia; but in cells exhibiting so much movement it is very easy to transfer what is seen in one view to a sketch representing in general a different view, unless one is keenly on the lookout for such a point. Probably also many of our somewhat classic figures are not made from camera lucida drawings of quiescent individuals, as they should be to determine such features.

In the asexual reproduction of *C. caudata*, Wille describes the division as longitudinal, after the disappearance of the cilia of the mother cell, resulting in the formation of four zoospores. On several occasions when I examined material with my travel microscope in the field in late afternoon, or freshly collected material in early evening, I found numerous individuals which contained the four fully formed zoospores while the mother cell was still swimming actively. It was only in the more rarely seen divisions of material kept for a day or two after collection that I found non-motille cells in process of reproduction; such cells may have lost the power of movement earlier because of poorer aeration than in the normal habitat. This division of actively motile cells I have found to be normal in several other species of *Chlamydomonas*. I was unable to find examples which would actually show more of the details of the process of division than indicated by Wille, who merely furnishes one figure of completed division. I did, however, observe that a large majority of cases showed two of the daughter cells with their anterior end directed forwards, and two with the anterior end directed backwards (FIG. 49). In other species of *Chlamydomonas* such as arrangement of the daughter cells is characterstically found when the first plane of division appears transverse to the longitudinal axis of the mother cell, or when, though fundamentally longitudinal, the plane becomes placed transversely by a rotation of the entire protoplast. During the past season I have had the opportunity of observing this latter method many times in a new species to be described in a future paper: this species furnished one exceptional case in which both divisions, carefully followed, were strictly longitudinal, resulting in four zoospores all headed toward the anterior end of the mother cell. This exceptional arrangement I also found in rare instances in *C. caudata* (FIG. 50). Wille's figures (**15**, *pl. 3, f. 8*) which shows the same arrangement, appears, therefore to be untypical, and I am obliged to believe that the first division of this species, though properly longitudinal, would regularly show as a transverse cleavage, in consequence of a rotation of the protoplast to such an extent that its anterior pole comes to lie in a position nearly in the middle of the length of the cell.

Reichenow (**9**, **10**, pp. 35–39) it is true, has expressed the view that the division in all species of *Chlamydomonas* may be fundamentally longitudinal, an apparent transverse division being al-

ways preceeded, in his opinion, by a rotation of the protoplast, so that he believes the manner of cell division to have no phylogenetic significance. Attractive as this view is in its possibility of simplifying the idea of direct transmission of polarity to the daughter cells (and my own recent studies on *Chlamydomonas* somewhat tend to support such a view) nevertheless we still lack sufficiently definite and accurate observations to permit the denial that an actual transverse division may occur in some species.

If in *C. caudata* the first cleavage plane is actually or by rotation of the protoplast transverse (as appears to me most probable), and if we accept the current view derived from Dill (4) that the most primitive species are those in which both divisions are longitudinal, then *C. caudata* could hardly be regarded as an actual transitional form between the ordinary ovoid *Chlamydomonas* type and the *Brachiomonas* group, since all species of the latter genus appear to have retained the primitive mode of division by two strictly longitudinal cleavages, in spite of their advancement in other respects. Nevertheless, in its general morphology, and in the similarity of its method of formation of aplanospores (rarely found in other species of *Chlamydomonas*), *C. caudata* must be very close to the ancestral line which has given rise to the four forms of *Brachiomonas* recognized above; and these present a rather close ascending series, with perhaps a definite orthogenetic tendency beginning with *B. simplex*, which is barely emerging from the *Chlamydomonas* stage, through *B. submarina f. obtusa* and then *B. submarina* (type) to *B. gracilis*, an extremely slender form, so far removed from any *Chlamydomonas* type as to justify thoroughly the retention of *Brachiomonas* as a genus.

In certain graphic schemes of the phylogeny of the chlamydomonads—e. g. those of Wille (**15**) and West (**14**, p. 163)—the genus *Lobomonas* Dangeard is indicated as representing an intermediate stage between *Chlamydomonas* and *Brachiomonas*; but the two known species of *Lobomonas*, as well as two new species to be described by the writer in the next number of this journal, are very minute forms, and in their general morphology exhibit by no means so good a connection as is shown by *C. caudata* and *B. simplex*. *Lobomonas* must be regarded as a special offshoot from the *Chlamydomonas* line not leading to anything higher, so far as we know at present.

BARNARD COLLEGE,
COLUMBIA UNIVERSITY

LITERATURE CITED

1. **Bohlin, K.** Zur Morphologie und Biologie einzelliger Algen. Öfvers. K. Vet. Akad. Förhand. **54**: 507–529. 1898.
2. **Chodat, R.** Algues vertes de la Suisse. Berne. 1902.
3. **Dangeard, P. A.** Mémoire sur les Chlamydomonadinées ou l'histoire d'une cellule. Le Botaniste **6**: 65–292. *f. 1–20.* 1899.
4. **Dill, O.** Die Gattung *Chlamydomonas* und ihre nächsten Verwandten. Jahrb. Wiss. Bot. **28**: 323–358. 1895.
5. **Harper, R. A.** The structure of protoplasm. Am. Jour. Bot. **6**: 273–300. 1919.
6. **Lagerheim, G.** Studien über artkische Cryptogamen. Tromsø Mus. Aarhefter **17**: 1–24. 1895.
7. **Pascher, A.** Über die Kreuzung einzelliger, haploider Organismen: *Chlamydomonas.* Ber. Deutsch. Bot. Ges. **34**: 228–246. *f. 1–5.* 1916.
8. **Printz, H.** Kristianatraktens Protococcoideer. Vid.-Selsk. Skrift. I. Mat. Naturv. Klasse **1913**[6]: 1–123. *pl. 1–7* + *f. 1, 2.* 1914.
9. **Reichenow, E.** Untersuchungen an *Haematococcus pluvialis* und eingen anderen Flagellaten. Sitz.-ber. Ges. Naturf. Freunde Berlin **1909**: 85–91. 1909.
10. ————. Untersuchungen an *Haematococcus pluvialis* nebst Bemerkungen über andere Flagellaten. Arb. Kais. Gesundheitsamte **33**: 1–45. *pl. 1, 2.* 1909.
11. **Thompson, D'Arcy W.** On growth and form. Cambridge. 1917.
12. **West, G. S.** Some critical green algae. 2 *Brachiomonas submarina.* Jour. Linn. Soc. Bot. **38**: 281–283. *pl. 20. f. 7–19.* 1908.
13. ————. Algological notes XX, XXI. Jour. Bot. **54**: 2–5. *f. 1, 2.* 1916.
14. ————. Cambridge Botanical Handbooks. Algae 1. Cambridge. 1916.
15. **Wille, N.** Algologische Notizen XI. Über die Gattung *Chlamydomonas.* Nyt Mag. Naturvid. **41**: 109–162. *pl. 3, f. 4–11.* 1903.

Description of plates 3 and 4

All drawings made with the aid of camera lucida from living material in hanging drop cultures (or in some cases freshly fixed with osmic acid vapor): PLATE 3 done with Leitz 1/12 in. oil immers. obj. and compens. oc. 6, magnifica-

tion about 1440 diameters; plate 4 done with Spencer 2 mm. oil immers. obj. and Leitz compens. oc. 6. magnification about 1300 diameters; both reduced in reproduction to about 720 diameters.

PLATE 3

FIGS. 1–7. BRACHIOMONAS SUBMARINA Bohlin, type form

FIG. 1. Vegetative cell; posterior horn filled with colorless cytoplasm.

FIG. 2. Chromatophore filling the extremities: Dec. 1919.

FIG. 3. Colorless cytoplasm in extremities; stigma and pyrenoid lying on the under side of the cell: March, 1921.

FIG. 4. Anterior polar view of similar cell; vacuoles in chromatophore.

FIG. 5. Slender form similar to Bohlin's *f. 1b* of *B. gracilis* (fixed with osmic acid vapor).

FIG. 6. Moderately slender form: showing typical position taken by living zoospore temporarily at rest, the pyrenoid and stigma lying in the center of the under side (osmic vapor): April 1921.

FIG. 7. Anterior polar view of similar cell, showing cilia stretching out in plane perpendicular to that in which the nucleus, pyrenoid, and stigma typically lie.

FIGS. 8–27. BRACHIOMONAS SUBMARINA forma OBTUSA Hazen

FIG. 8. Chromatophore contracted, arms and horn filled with colorless cytoplasm (osmic vapor): November, 1919.

FIGS. 9, 10. Arms empty, colorless cytoplasm extending into posterior horn (osmic vapor).

FIG. 11. Anterior polar view of similar cell, living.

FIG. 12. Arms and horn nearly empty (osmic vapor).

FIG. 13. Posterior polar view of similar cell, living, showing typical relative positions of organs as in FIG. 7; a large colorless cytoplasmic region around the nucleus.

FIG. 14. Older cell showing no trace of arms.

FIG. 15. Cell from old culture with only a suggestion of lateral arms; stigma and pyrenoid lying underneath the nucleus.

FIG. 16. Senescent cell with contracted protoplast, preparing to form an aplanospore: December, 1919.

FIG. 17. Young cell showing nucleus posterior to pyrenoid.

FIG. 18. Posterior polar view of similar cell.

FIG. 19. First division of motile cell; 10:30 P. M. (osmic vapor).

FIG. 20. Eight zoospores fully formed in motile mother cell.

FIG. 21. Four zoospores: cilia of mother cell no longer present.

FIGS. 22–27. Successive stages in division resulting in eight zoospores (November, 1919): FIG. 22 at 2:40 P. M.; FIG. 23 at 3:40; FIG. 24 at 4:50; FIG. 25 at 5:30; FIG. 26 at 6:20; FIG. 27 at 7:40. The zoospores escaped before 8:40.

PLATE 4

BRACHIOMONAS SIMPLEX Hazen

Drawn 29 June—4 July, 1920, Aalesund, Norway.

FIGS. 28, 29. Typical mature vegetative cells: in FIG. 29 the stigma lies slightly underneath.

FIG. 30. Anterior polar view of similar cell.

FIG. 31. Cell on which the "bumps" representing lateral arms are undeveloped.

FIG. 32. Anterior polar view of similar cell.

FIG. 33. Protoplast ertracted from one of the "bumps" (osmic vapor).

FIG. 34. Large older cell with more posterior pyrenoid.

FIG. 35 Eight zoospores in motile cell; 12:45 A. M.

FIG. 36. Four zoospores in motile cell.

FIG. 37. Young zoospore free; chromatophore filling posterior horn.

FIG. 38. Three of the four zoospores have escaped through a triangular rent in the wall of the mother cell.

FIG. 39 Similar to FIG. 31, but with more typical position of pyrenoid (osmic vapor)

FIG. 40. Thirty-two gametes in motile mother cell; 10 A M.

FIGS. 41, 42. Gametes at beginning of conjugation (osmic vapor).

FIG. 43. Motile zygote, momentarily resting; nuclei not fused.

FIG. 44. Zygote, cilia having disappeared, nuclei fused; two stigmata still present.

FIG. 45. Zygospore twenty-four hours after conjugation.

In the living cells the pyrenoid actually appears upon casual observation to be green because imbedded in the green chromatophore. It is believed, however, that the printing of the pyrenoids in the same sepia as the cell-wall and protoplasmic outlines represents more truthfully their real nature.

Note. As final proof of this paper is received, the writer has just made a mid-April excursion to the Massachusetts coast, where he has collected typical *Brachiomonas submarina* Bohlin from Cape Ann to Buzzards Bay, at Rockport, Gloucester, Marblehead, Nahant, and Fair Haven. There has been no indication of other species of *Brachiomonas* or of *Chlamydomonas caudata* at these stations.



1-7. BRACHIOMONAS SUBMARINA BOHLIN
8-27. BRACHIOMONAS SUBMARINA FORMA OBTUSA HAZEN



BRACHIOMONAS SIMPLEX HAZEN

New records and other notes on North Dakota plants*

O. A. Stevens

The present paper is supplementary to Bergman's Flora of North Dakota (in Sixth Bien. Rept. Agr. Coll. Survey [1911–1912]. 1918) and, as such, will follow the same arrangement and nomenclature. The purpose is to record additional species, designated by a dagger (†), and other notes of special interest. During the years 1910 to 1920 I have had opportunity to visit nearly every part of the state and, incidental to economic work, to obtain many specimens of interest. Some of these were noted in the Flora and others have been reported by Lunell (Am. Mid. Nat. **4–6**. 1917–1919). I have also included in the paper a few corrections which have come to notice and revised certain groups in which I have become especially interested. Where not otherwise stated, the specimens listed have been collected by myself.

Lepidium ramosissimum A. Nels. I do not know why Bergman reduced this to a synonym of *L. densiflorum* Schrad. The two species are very different and may be distinguished as follows:

L. densiflorum. Annual, stem usually simple below and with several long racemes above; basal and lower stem leaves serrate; pods distinctly margined and broadest toward the apex.

L. ramosissimum. Biennial, bushy branched from the base; basal and lower stem leaves pinnatifid, the lobes linear or oblanceolate; pods not margined above, elliptic in outline.

The following specimens belong to *L. ramosissimum*: Tower City, *Bergman 870*; Valley City, Aug. 13, 1912, (first year rosettes); Courtenay, June 20, 1911; Kensal, July 26, 1915, (rosettes, also flowering stems); Manfred, July 12, 1913; Sykeston, July 15, 1891, *Bolley*; Sheyenne, July 4, 1908, *Lunell;* Leeds, Aug, 1, 1912; Bathgate, July 22, 1892, *Lee;* Ganbetta; *Bell 264*; Rugby, *Bergman 2611*. This species is quite common throughout the central part of the state, growing about streets and dooryards rather than in fields. At Langdon on July 19, 1918, I noticed the rosettes of *L. ramoisssimum* fairly covering

* Contribution from Department of Botany, North Dakota Agricultural College and Experiment Station.

the ground. In a dooryard at Courtenay the species grew mixed with *L. densiflorum*. The latter had racemes about 3 cm. long with numerous well-developed pods, while the plants of *L. ramosissimum* had only a few half-developed pods.

† LEPIDIUM DRABA L. Several years ago (Ann. Rept. North Dakota Exp. Sta. **22**: 80. 1912) I called attention to the possibility of this being introduced in Turkestan alfalfa seed. On June 6, 1918, a flowering plant was sent me by Mr. H. A. Nelson of Ray. [A well established colony has since been found along the railroad track at Moorhead, Minnesota, May 22, 1921.]

† SISYMBRIUM LOESELII L. Determination verified by Paul C. Standley and specimens deposited in herbaria of the U. S. National Museum and the New York Botanical Garden.

Annual, 6–9 dm. high, sparingly hirsute with simple flattened hairs, these less abundant above but present on upper stems and pedicels; leaves runcinate, the terminal segment hastate and irregularly denticulate, lower stem leaves sometimes 1 dm. long; racemes becoming 3–4 dm. long, pedicels widely spreading one third to one half the length of the slender pods, which are ascending and about 3 cm. long with three-nerved valves.

Devil's Lake, July 15, 1920 (common about the streets); Tappen, July 12, 1919 (a single plant in the street; several in an old field of *Bromus inermis* Leyss. several years before). The description was drawn from the growing plants and specimens collected at Devil's Lake. The plant has somewhat the aspect of *Sophia intermedia* Rydb., but the leaves are quite different and the racemes are fewer but longer and coarser. The relative height of flowers and young pods mentioned by Koch and others seems scarcely distinctive. In this plant the pods do not surpass the flowers in normally developed racemes, but examples are frequent where they do so, apparently as a result of arrested development.

ERUCASTRUM POLLICHII Schimp. & Spenn. Park River, Aug. 8, 1913; Williston, Aug. 15, 1915; Oakes, July 18, 1919; Cooperstown, Aug. 25. 1919. This has now been found at many places, chiefly along the railroad tracks. The seeds have been identified in several samples of timothy and millet coming from near Grand Forks or a short distance north or south of this locality. The pods are not flattened, as stated in Bergman's key, but rounded, the valves with a rather prominent mid-nerve; racemes leafy bracted. The plants grow vigorously in late fall as shown by a flowering specimen collected at Fargo, Oct. 26, 1914.

MALVA VERTICILLATA L. I have not seen the specimen which Bergman referred to this species but am inclined to think that it was *M. borealis* Wallm., which is the common plant at Fargo and other places in the state. A specimen of what seems to be *M. crispa* L. † was sent by A. G. Sorlie, Grand Forks, Oct. 10, 1916.

EUPHORBIA ESULA L. Wahpeton, June 3, 1920, *Geo. P. Wolf;* Hazelton, July 15, 1919, in field of *Bromus inermis*; New Rockford, July, 1911, *J. R. Campbell*; Mona, June 5, 1914, *Albert Graves*; Ray, July 2, 1917, *H. A. Nelson.* In one field near Fargo patches are scattered over 20 acres. Apparently a bad weed, spreading by the roots. Time and manner of introduction unknown.

† SILENE DICHOTOMA Ehrh. Edgeley, July, 25, 1917, *Brenckle;* Langdon, July 19, 1918; Regent, July 20, 1916, *A. C. Goldtrap.* In timothy fields. Plants grown from seed proved to be coarse biennials larger than *S. noctiflora.*

† SILENE FABARIA (L.) Sibth. & Sm. Specimens determined by Paul C. Standley.

Biennial; glaucous, much branched, 6–9 dm. high. Basal and lowest stem leaves narrowed at the base, larger stem leaves ovate, sessile; flowering stems dichotomous with a flower in the fork, each branch with several remote clusters of three to seven flowers each. Flowers about 1 cm. wide, white, opening at night; petals cleft to the throat, not appendaged, the lobes oblanceolate, rounded at the apex; calyx indistinctly fifteen-ribbed, closely investing the ovate horny capsule, but not especially enlarged in fruit.

Venturia, July 16, 1919; Oakes, July 18, 1919. Lower stems and leaves with very much the aspect of *Vaccaria Vaccaria.* A specimen in the herbarium of Dr. J. F. Brenckle was collected at Kulm by him in 1916, and he states that he had found the plant there previous to that date. Apparently well established at Oakes and Venturia along roadsides. Dr. Standley writes that this is probably the first time it has been collected in this country.

† LYCHNIS ALBA Mill. Devil's Lake, July 13, 1920.

CHENOPODIUM

For several years I have been interested in *C. album* and related forms, especially in regard to seed characters. The plants are rather difficult to collect as the seeds are slow in maturing and the lower leaves are likely to be lost by the time the fruiting branches are well matured. With respect to the adherence of the pericarp it is to be noted that it rubs off more easily in fresh material.

In the fall of 1916 I collected twenty-six specimens in the vicinity of Fargo and submitted them to Dr. Standley, who determined them as follows:

C. album (4);

† *C. paganum* Reichenb. (11);

C. paganum, approaching *C. album* (8);

† *C. ferulatum* Lunell (3).

Apparently *C. paganum* is the commonest form here. I have not as yet been able to make much further progress in clearing up the relations of these forms but wish to offer what few notes I have.

C. ferulatum (which Standley notes is perhaps not sufficiently distinct from *C. album*) is separated by the fruiting calyx being open, exposing the fruit. While this does not seem to be quite constantly true for *C. ferulatum*, I find it also in a rather variable degree in the *C. album* specimens, being most prominent in well-matured plants. No. 11, determined as *C. paganum*, has calyces closed in specimen collected Aug. 25, but fruit from same plant collected in September shows them quite open. As to *C. paganum*, most of the plants were either so badly infested by an insect or so late in maturing that very little mature seed was obtained. From material at hand I would describe the seeds as follows:

C. album—1.2–1.4 mm. wide, upper surface flattened and with shallow, oblong pits (which show through the pericarp).

C. paganum—1.2–1.4 mm. wide, upper surface dull, neither flattened nor pitted.

C. ferulatum—1.2–1.4 mm. wide, similar to *C. paganum*.

C. leptophlyllum—0.9 mm. wide, not flattened, smooth and shining.

The size of the seed of *C. paganum* is given by Standley (N. Am. Fl. **21**: 21. 1916) as 1.3–2 mm. In one of the twenty-six specimens, a plant collected near the woods along the Red River, the seeds are variable in size, measuring 1.4–2 mm., but in all the others they are scarcely larger than those of *C. album*.

The three specimens of *C. ferulatum* differ among themselves. One, collected in an old garden (No. 3), which I take to be typical, is pale grayish green, simple below and with short branches above, the leaves oblong-rhombic with four or five short teeth on each side; another (No. 9, labelled by Standley as apparently a form of *C. ferulatum*) is bushy branched with narrow entire leaves. The third (No. 2) is in a rather advanced condition with lower leaves mostly fallen; the flowering branches

are unusually thick and dense and the seeds are pitted on the upper side. I am inclined to think it is a plant of *C. album* with an unusual amount of well-matured fruits and therefore with the spreading calyces unusually prominent.

The seeds of these species are described by Standley as "nearly smooth, black and shining" except in *C. ferulatum*, where they are said to be "puncticulate," but I find in our material that *C. album* has pitted seeds and *C. ferulatum* smooth (excepting as in No. 2 just noted; type material from Lunell also examined). The seeds of *C. leptophyllum* seem to be quite easily distinguished from those of the other common field species by their smaller size.

† CHENOPODIUM PRATERICOLA Rydb. Narrows, Aug. 12, 1913, Lunell.

† ATRIPLEX PATULA L. Wahpeton, Aug. 6, 1919. Along a street by buildings. Dr. Standley, who examined a specimen, writes that this species is perhaps only a form of *A. hastata* L., but we seem to have no other specimens which would be referred to this form.

† SUCKLEYA SUCKLEYANA (Torr.) Rydb. Belfield July, 4, 1914.

CORISPERMUM

Following Standley's treatment of this genus (N. Am. Fl. 21: 79–80. 1916) I would place our material as follows:

C. hyssopifolium L. Mandan, *Wright 610.*

† *C. nitidum* Kit. Sandhills near Anselm, Aug. 21, 1918.

† *C. villosum* Rydb. Sandhills near Anselm, Aug. 21, 1918; also all other specimens referred by Bergman to *C. hyssopifolium*, except *Wright 610.* The Fargo specimen of *C. villosum* was collected along the railroad (doubtless introduced in gravel) and no plants have been observed since. It was a well matured plant with very little pubescence except on the bracts. The Anselm plants are young, upright, rather slender, and densely pubescent.

POLYGONUM HYDROPIPER L. Hankinson, July 30, 1919. Lunell (Am. Mid. Nat. **5**: 184. 1918) has referred the Ft. Ransom specimen to *P. punctatum leptostachyum* (Meisn.) Small, but it has dull akenes and flowers in the lower axils, as in *P. Hydropiper.*

ASCLEPIAS SYRIACA L. Westfield, July 16, 1919. This does not seem to be common so far west. The specimen reported

by Bergman from Morton County (*Bell 30*) is undoubtedly *A. speciosa* Torr.; it is sterile but has the characteristic pointed leaves of that species. The specimen reported by Bergman as possibly a hybrid of the two species seems scarcely different from *A. syriaca.*

† ASCLEPIAS PUMILA (A. Gray) Vail. Mr. W. W. Eggleston has examined our material of *A. verticillata* L. and refers the following to *A. pumila:* Esther, *Bell 542*; Fleak, *Bell 1370.*

† ACERATES LANUGINOSA (Nutt.) DC. Janesburg, *Bell 30*, of Bergman's catalogue should be referred to this species instead of to *A. viridiflora* Ell.

† VERONICA MARITIMA L. Kongsberg, Sept. 1918, *Fred Schmidt, Jr.* A few plants in a grove of planted trees. Determined by F. W. Pennell.

CUSCUTA ARVENSIS Beyr. The Logan County record is by Brenckle, Aug. 18, 1912. We have also the following records: McLeod, *Bell 385*: Valley City, July 26, 1913; Enderlin, Aug. 20, 1918. Dr. T. G. Yuncker, who has examined portions of the Logan County and Valley City specimens, writes that he would call them C. *pentagona* Engelm. The other specimens seem to be the form which he calls *C. calycina* Engelm. The former were growing upon plants of the dry prairie, while the latter preferred those of riverbanks and similar localities.

CUSCUTA GRONOVII Willd. Fargo, Aug. 14, 1890, *Waldron*, is the only specimen which I consider typical. When preparing my paper on *Cuscuta* (Am. Jour. Bot. **3**: 185–188. 1916) I was unable to find fresh material, but all that I have found at Fargo since that time agrees better with what I had called *C. plattensis* A. Nels. The habitat of *C. plattensis*, as stated by Nelson, hardly indicates this plant and I become doubtful of its identity. Dr. Yuncker writes me that an examination of the type of *C. plattensis* shows my plant to be quite different, although the description of the species as given by Nelson agrees. He refers my specimens to *C. Gronovii* var. *curta* Engelm. In a flax field at Ft. Ransom, where considerable damage was caused, this form and *C. Coryli* Engelm. were both present, sometimes separate and at others tangled together on the same host plant. The field was a small one next to the woods of the river.

PHLOX

Following the treatment of the western forms by E Nelson (Ninth Rept. Wyoming Agr. College, 1899), our plants would seem to be separated more satisfactorily as follows:

Leaves 5–10 mm. long; flowering stems one-flowered. *P. Hoodii.*
Leaves 1–2 cm. long; flowering stems two-to several- flowered: more upright, the bark peeling off in shreds. *P. andicola.*

PHLOX ANDICOLA (Britton) E. Nelson. *P. Douglasii* of Bergman's Flora, in part. This seems to be better separated from *P. Hoodii* Rich. by the above characters than by the size of the flowers. The Medora and Washburn specimens referred by Bergman to *P. Douglasii* Hook. are evidently *P. Hoodii.*

† LAPPULA CENCHRUSOIDES A. Nels. Marmarth, July 4, 1918, in sandy soil of river valley. Habit of growth similar to *L. occidentalis* (Wats.) Greene, nutlets similar to those of *L. Lappula* (L.) Karst. but larger, the spines longer, and with a row of elongated tubercles on the middle of the back of the nutlets. The following, previously referred to *L. Lappula*, also belong here: Medora, *Bergman 1276*; same locality, June 19, 1910; Williston, *Bell 25*.

† LAPPULA CUPULATA FOLIOSA (A. Nels.) Nels. & Macbr. Marmarth, same as preceding. Often three of the nutlets of a flower have the spines confluent, forming a spreading border, the fourth nutlet with simple spines. In the specimens collected, however, there seems a decided tendency for the spines to be simple on all nutlets of the lower flowers. There are also nutlets with spines of intermediate form, broadened but not united. Counts on three plants show nutlets as follows:

Plant No. 1—80 united, 71 simple, 40 intermediate;
Plant No. 2—61 united, 106 simple, 34 intermediate;
Plant No. 3—47 united, 47 simple, 14 intermediate.

The color of the corolla is an uncertain character, white flowers often appearing bluish in the dried specimens.

† LAPPULA TEXANA HOMOSPERMA (A. Nels.) Nels. & Macbr. Marmarth, same as preceding, occasional plants in patches of *L. occidentalis*, all plants dried up and only fruits collected. From a planting of fruits the following season about twenty-five plants of *L. occidentalis* were obtained from fruits of that species. No plants were secured from the few fruits of the other two species.

† LAPPULA FLORIBUNDA (Lehm.) Greene. Lake Ibsen (Leeds) July 4 and 17, 1914, *Lunell*; Pleasant Lake, June 29, 1920.

† CRYPTANTHA CALYCOSA (Torr.) Rydb. Bowman, June 23, 1918, in loose burned clay around large boulders of the same material.

AMSINKIA MENZIESII (Lehm.) Nels. & Macbr. *A. intermedia* F. & M. Rugby, July 7, 1917, and July 27, 1918, a number of plants along the railroad track near the station; the first collection in flower, the second in fruit. Bergman's specimen from Pembina is evidently this species also, instead of *A. lycopsoides* Lehm.

† LAMIUM AMPLEXICAULE L. Langdon, July 18, 1918, a quantity in a dooryard.

† DRACOCEPHALUM THYMIFLORUM L. Belfield, July 4, 1914, several plants in a field of *Bromus inermis*. Several other introduced plants were also in the field: *Potentilla argentea* L., *Campanula sibirica* L., *Chrysanthemum Leucanthemum* L. and *Bromus tectorum* L. To be consistent with Bergman's nomenclature this plant should perhaps be referred to *Moldavica* but I have not thought it desirable to make such a change at this time. The plant is different in appearance from *D. parviflorum* Nutt., being slender and having very small flowers.

† LYCOPUS COMMUNIS Bicknell. Pleasant Lake, July 26, 1912, *Lunell*; Anselm, Aug. 29, 1920 (common in alder swamps). The Anselm plant has been verified by P. A. Rydberg.

† SALVIA LANCEOLATA Willd. In fields. Carrington, Aug. 27, 1919; Mandan, Sept. 13, 1920.

CHAMAERHODOS ERECTA (L). Bunge. This was included in the seventh edition of Gray's Manual, the record being apparently based on a specimen collected near Crookston, Minn. (Minnesota Bot. Studies **2**: 584. 1901). I had thought that it might have been introduced in gravel as railroad ballast, but so far as I have been able to learn the locality mentioned is a gravel pit. It is probably one of the instances of a plains species occuring on the eastern edge of the Red River Valley. I have not seen the plant east of the hills along the Sheyenne River at Sheyenne and Valley City.

† CRATAEGUS MOLLIS T. & G. Fargo, May 26, 1917. One tree is in a thicket east of the fair grounds, and quite a number occur in a bend of the river two miles farther north.

† MEDICAGO FALCATA L. Oakes, July 18, 1919, a plant along the railroad tracks.

† ASTRAGALUS PARVIFLORUS (Pursh) MacM. Marmarth, July 3, 1918. Quite common on the higher parts of the hills.

† HEDYSARUM sp. Sentinel Butte, Aug. 0916, *Brenckle*, a single pod collected. Reported by Arthur (N. Am. Fl. **7**: 450. 1921) as *H. cinerascens* Rydb.

† EVONYMUS ATROPURPUREUS Jacq. Owego, Sept. 1916, *R. A. Shunk*.

† LOMATIUM MACROCARPUM (Nutt.) Cov. & Rose. Dunseith, May 18, 1918, *Lunell*, and fruits collected at Minot in June, 1910, seem properly referred to this species.

† SICYOS ANGULATUS L. I have seen a specimen collected by A. H. Shunk along the Sheyenne River near Anselm.

† CAMPANULA SIBERICA L. Belfield, July 4, 1914, a single plant found in a field of *Bromus inermis*.

† XANTHIUM PENNSYLVANICUM Wallr. A specimen in the Gray Herbarium from Leeds, Aug. 21, 1902, *Lunell*, is referred by Millspaugh and Sherff (Field Mus. Nat. Hist. Bot. Ser. **4**: 33. 1919) to this species. Some specimens of *Xanthium* collected along the river at Fargo in 1919, to show variations in burs, were determined by Sherff as *X. italicum* Mor. (*X. canadense* of the Flora) and *X. acerosum* Greene, † with the comment that the latter was perhaps not distinct.

† CHRYSOTHAMNUS NAUSEOSUS (Pall.) Britton. Williston, Aug. 11, 1915, on the hills along the Missouri River about fifteen miles southeast of the city. A plant of quite different appearance from *C. graveolens* (Nutt.) Greene, which is common there and on the buttes in the bad lands—the crown low, and with gray branches only 1–2 dm. long. This is the plant referred by Lunell (Am. Mid. Nat. **5**: 41. 1918) at my suggestion to *C. formosus* Greene, but it evidently is not that species.

HELIANTHUS GIGANTEUS L. The form referred to in Bergman's Flora under this name is quite common in the central (and western?) part of the state, but I am as yet in doubt as to its status. It is evidently Lunell's *H. nitidus* (Am. Mid. Nat. **1**: 235. 1914). Specimens from Valley City and New Rockford were determined by Standley as *H. tuberosus*. The Fargo specimens are *H. Maximiliani* Schrad.

HELIANTHUS GROSSE-SERRATUS Martens. The Fargo specimen of the Flora is certainly *H. Maximiliani*. The stem is only slightly hispid above and the leaves are broader and more nearly flat than usual. The Kenmare specimen (*Bergman 2744*) belongs to the preceding form.

† MADIA GLOMERATA Hook. Spring Brook, Aug. 17, 1915, a quantity in a prairie slough near the town.

† ARTEMISIA PABULARIS (A. Nels.) Rydb. Mandan, Sept. 11, 1020. Determined by Rydberg. There seems to be no other specimen among our material which resembles this. Mr. Thysell of the Northern Plains Station showed me the plants, only two places where they were growing being known.

† SENECIO MANITOBENSIS Greenman. Bottineau, July 8, 1917. Determined by J. Lunell. In the meadow of a dried up pond in the Turtle Mts.; Towner, July 13, 1911, *Lunell* (specimen in the Gray Herbarium).

† CENTAUREA PICRIS Pall. Spring Brook, Aug. 11, 1918. Determined by Standley. Received from Jacob Widman who referred to it as abundant in a field. Especially a few years ago the akenes were often found in Turkestan alfalfa seed, but this is apparently the first record of its having become established.

SONCHUS ARVENSIS L. Ellendale, July 24, 1919. Several spots around buildings in the town.

†SONCHUS ULIGINOSUS Bieb. *S. arvensis* of Bergman's Flora, Fargo, Aug. 1, 1916; Oakes, July 18, 1919; Crosby, June 7, 1919, *J. H. Phelps;* Williston, Aug. 1915. Specimens of this and of the preceding were examined by Standley, who stated that no North American specimens were in the U. S. National Museum collections. A specimen sent to the New York Botanical Garden was reported by Rydberg as apparently new to this country*. This seems rather strange in view of the fact that it is the common form through the Red River Valley, and westward in the northern part of North Dakota to somewhat beyond Devil's Lake, southern Manitoba, and western Minnesota. It is said to have first appeared at Portage Plains, Manitoba, about 1900.

This form differs from *S. arvensis* in the absence of glandular hairs on the upper stems and involucres. While I have not been able to compare carefully fresh material, I believe the heads are smaller, paler, and the rays more inclined to become recurved. Vegetative reproduction is by long horizontal roots and not rootstocks as commonly stated. The development of the terminal head is often stopped before flowering and several flowering branches arise just below it.

* Small has recently listed the species as an addition to the American flora, from Pennsylvania, where it was collected in 1921 (see Torreya **21**: 100. 1922).

LACTUCA VIROSA L. Many European authors describe this as having horizontal leaves and black, broadly margined akenes. This does not apply to our plants. The pinnatifid-leaved form is only occasional in the state (Kulm, *Brenckle 782*; Bottineau, July 23, 1918; Fargo, Aug. 17, 1920). Pammel's notes (Rhodora 20: 180–181. 1918) on the replacement of the entire-leaved form by the pinnatifid in Iowa suggest that it will be interesting to watch for a similar case here.

† CREPIS OCCIDENTALIS Nutt. Bowman, June 23, 1918. Quite common on Twin Buttes.

† CREPIS CAPILLARIS Roth. Fargo, Aug. 26, 1920; Willow City, *F. M. Rich.*

† SPIRODELA POLYRHIZA (L). Schleid. Fargo, Aug. 30, 1919, a few plants among masses of *Lemna*.

† CYPERUS DIANDRUS Torr. Anselm, Aug. 1916, *R. A. Shunk.*

† CYPERUS ESCULENTUS L. Fargo, Oct. 2, 1920.

SCIRPUS VALIDUS Vahl. I cannot support Bergman's separation of most of our material as *S. occidentalis* (Wats.) Chase. There do seem to be two forms present, one with spikelets as figured in Gray's Manual (ed. 7, *f. 290*), the other with more rigid panicles and longer spikelets. The specimens, however, do not match Bergman's separation. Both of these forms were found in a small patch at Oakes on July 18, 1919, the first occupying a definite part of the patch. Specimens of each were identified by Agnes Chase as *S. validus*.

CAREX OBTUSATA Liljebe. Verified by Standley. Fargo, June 24, 1920. Quite abundant, at least in one place in low prairie. Kensal, *Bergman, 1744*, is alsothis in stead of *C. stenophylla* Wahlenb.

The following additional species of *Carex* were recorded by Lunell (Am. Mid. Nat. 3: 234–237. 1916) from specimens determined by Mackenzie: *C. Hookeriana* Dewey, *C. athrostachya* Olney, *C. tenera* Dewey, *C. praticola* Rydb., *C. Emoryi* Dewey, *C. scirpiformis* Mackenzie, *C. Parryana* Dewey, *C. laeviconica* Dewey and *C. atherodes* Spreng.

† SYNTHERISMA SANGUINALE (L.) Dulac. Fargo, Oct. 2, 1920.

† PANICUM PERLONGUM Nash. Verified by Hitchcock. Fargo, June 24, 1920. Very abundant in one place in low prairie.

† SPOROBOLUS ASPER (Michx.) Kunth. Mayville, Aug. 21, 1919: Fargo, Aug. 16, 1920; Steele, Sept. 10, 1920.

FESTUCA VIRIDULA Vasey. Lunell (Am. Mid. Nat. 4: 224 1917) has reported this species from Dunseith. I have examined carefully a specimen received from him ("det. by U. S. Dept. Agr.") but can see no reason why it should not be referred to *F. Hallii* (Vasey) Piper.

BROMUS JAPONICUS Thunb. Fargo, Oct. 1918. Determined by Agnes Chase. All of the specimens referred by Bergman to *B. commutatus* Schrad., and several others not reported, seem to belong here.

† LOLIUM RIGIDUM DUTHIEI Hook. Determined by Agnes Chase.

Annual: culms 3 to 4 dm. high, rather stiff, somewhat scabrous above; leaves upright, 5 mm. wide, 1 dm. or more long, glabrous or nearly so; glume about three-fourths as long as the well developed spikelets, equalling the younger or shorter ones; spikelets five- to seven-flowered, the larger 2 cm. long; lemmas 8 to 10 mm. long bearing an awn 7 to 12 mm. long.

Described from a specimen collected July 7, 1919, grown from seeds found in a sample of wheat from the northeastern part of the state. An early maturing annual, not so large as *L. temulentum* L. and quite different in the longer, lanceolate, awned florets. The culms commonly bear a branch from the first node, some of these in the specimen cited being only 1 dm. long with spikes barely protruding. The Milton specimen cited by Bergman under *L. temulentum* belongs here, both it and plants from the 1919 culture having been examined by Mrs. Chase. The plant seems well established in that vicinity as evidenced by these cases, by material received at one or two other times and by seeds found in several samples of wheat.

† LOLIUM REMOTUM Schrank. In flax plots, Mandan, 1918.

AGROPYRON

Three species of this genus are of great economic importance in North Dakota. Having had occasion to pay particular attention to these, I find that as a rule, descriptions and figures seem to have been made from immature material, causing certain characters to be overlooked. The spikelets are well figured by Hillman (Bur. Pl. Ind., U. S. Dept. Agr. Circ. 73. 1911). I offer the following descriptions:

A. repens (L.) Beauv. Spikelets distinctly articulated to the rachis (so that they break away readily, with a rounded base), and with a strong transverse impression about 1 mm. above the base. Glumes from half as long to nearly equalling the spikelet, strongly nerved, with about a dozen short teeth on the keel* near the apex which is acute or obtuse, often bearing an awn 2–10 mm. long; margins of the glumes thin and translucent except toward the base where they are indurated, rounded, usually slightly separated exteriorly and quite widely interiorly.

A. Smithii Rydb. Spikelets not articulated (breaking away with a rough base) and only slightly impressed. Glumes about one-half as long as the spikelets, indistinctly nerved and without distinct teeth on the back; margins thin on lower third (meeting exteriorly), abruptly narrowed to a long stiff, acuminate point.

A. tenerum Vasey. Spikelets not articulated and only slightly impressed. Glumes about equalling the spikelets, acute, strongly nerved, the nerves all rough with small teeth, margins nearly straight.

† AGROPYRON DASYSTACHUM (Hook.) Scribn. A specimen from Bottineau, July 7, 1920, is referred here ("form which has been called *subvillosum*") by Hitchcock. Bowman, June, 30 1918, and Kenmare, July 15, 1913, are the same, and I think all the specimens previously referred to *A. molle* (excepting perhaps *Bell 385*) belong here.

† DRYOPTERIS THELYPTERIS (L.) A. Gray. Anselm, *R. A. Shunk.*

† DRYOPTERIS SPINULOSA (Retz.) Kuntze. Another specimen of the same collection as the preceding seems to belong here, as does also Walhalla, *Bergman 2000.*

Since the above was written I have received from Mr. F. P. Metcalf a reprint of an article (Jour. Washington Acad. Sci. **10**: 188–198. 1920) in which he reports eighteen additional species for the state. Several of the records suggest the probability of an error of indentification or of a difference in interpretation. In reply to my inquiry regarding a few of them Dr. Standley wrote that he has been able to find but one of the specimens, *Rumex Britannica* L. (McLean County), and that that seemed to be correctly determined.

* This is not actually the central nerve, the broader part of the glume being exterior and only one or two nerves on the side next the rachis.

Three new species of Cuscuta from Mexico

T. G. YUNCKER

(WITH THREE TEXT FIGURES)

In studying collections of *Cuscuta* from Mexico three species have been discovered which, it is believed, have not been hitherto described. Descriptions of these new species, together with camera lucida sketches, are given below. With the addition of these the number of species of *Cuscuta* now known to occur in Mexico is thirty-six.

Cuscuta dentatasquamata sp. nov.

Stems medium to slender. Flowers reddish, 3–4 mm. long, on pedicels shorter than or mostly about equal to the flowers, in rather compact, cymose clusters; calyx deep, lobes exceeding the corolla tube, triangular, acute, slightly overlapping at the base, thickened in the median region, forming a small carina on most of the lobes; corolla campanulate, lobes triangular, acute, upright or spreading, shorter than the tube and with slightly uneven margins; stamens shorter than the corolla lobes, filaments stoutish and about equal to the oval or orbicular anthers; scales about reaching the stamens, dentate, with



FIG. 1. CUSCUTA DENTATASQUAMATA Yuncker
A. Flower, × 5. B. Opened corolla, × 5. C. Individual scale, × 10. D. Opened calyx, × 5. E. Capsule, × 5.

few processes, bridged at about the middle, not firmly adherent to the corolla; styles slender, about equal to, or longer than, the depressed-globose ovary. Capsule large, depressed-globose, thin, somewhat irregularly circumscissile, carrying the withered corolla about it; seeds about 1.5 mm. long, globose, two to four in each capsule, hilum oblong, transverse.

This species falls in section EUGRAMMICA, subsection LEPTILOBAE, near *Cuscuta Choisiana* Yuncker. It differs from that

species, however, in the color of the flowers, and in the shape of the scales and of the calyx and corolla lobes. It superficially resembles *Cuscuta partita* Choisy, but differs from that species in possessing shorter corolla lobes, dentate scales, shorter filaments and larger flowers.

The only specimen examined was from Los Pinitos, Sonora, Mexico (*Hartman 119*), the type, in the Gray Herbarium.

Cuscuta cozumeliensis sp. nov.

Stems rather coarse. Flowers 4–5 mm. long, on pedicels about as long as, or shorter than, the flowers, in cymose clusters; calyx shorter than the corolla tube, lobes ovate-orbicular, overlapping, somewhat fleshy toward the base, margins very uneven; corolla campanulate, lobes much shorter than the tube, erect or spreading, overlapping, obtuse or rarely slightly acutish, margins uneven; stamens shorter than the lobes, filaments stoutish, about equal to, or slightly longer than, the oval anthers; scales not reaching the stamens, oblong, moderately fringed



FIG. 2. CUSCUTA COZUMELIENSIS Yuncker

A. Flower, × 5. B. Opened corolla, × 5. C. Opened calyx, × 5. D. Individual scale, × 10. E. Ovary, × 5.

with medium length processes, bridged at about a quarter of their height; styles slender, much longer than the depressed-globose ovary. The only specimen seen of this species is fragmentary and not ripe enough to show capsules. The ovary, however, gives indications that the capsule would be definitely circumscissile and probably carry the withered corolla about it.

This species falls in section EUGRAMMICA, subsection OBTUSILOBAE, near *Cuscuta macrocephala* Schaffner, from which it differs in the stamens not being sessile as they are in that species, and in having smaller stigmas and longer processes on the scales.

It might be confused with *Cuscuta floribunda* H. B. K., from which it differs in having shorter corolla lobes and shorter scales; with *Cuscuta corymbosa grandiflora* Engelm., from which it differs in the corolla not bulging between the filament attachments and with broader scales; and with *Cuscuta tinctoria* wartius, from which it differs in its larger flowers and proportionately shorter corolla lobes and shorter scales.

The specimen examined was from Cozumel Island, Yucatan, Mexico (*Gaumer 90*), the type, in the Gray Herbarium.

Cuscuta durangana sp. nov.

Stems slender. Flowers membranous, about 2 mm. long, on pedicels mostly as long as, or longer than, the flowers, in cymose panicles; calyx about as long as the corolla or scarcely reaching the sinus; lobes ovate, obtuse or, rarely, acutish, not overlapping; corolla campanulate; lobes ovate, obtuse, spreading, later becoming reflexed, about as long as the tube; stamens shorter than the lobes, the large, oval, versatile anthers about equal to the stoutish filaments; scales about reaching the stamens, obovate or spatulate, fringed about the upper half, bridged low; styles slender and longer than the globose ovary. Capsule depressed-globose, circumscissile, carrying the withered corolla about it; apparently mostly two-seeded; seeds ovate, slightly rostrate, about 1.5 mm. long, hilum oblong, oblique.



FIG. 3. CUSCUTA DURANGANA Yuncker A. Flower, × 5 B. Opened corolla, × 5 C. Opened calyx, × 5. D. Capsule, × 5. E. Individual scale, × 10.

This species falsl in section EUGRAMMICA, subsection OBTUSILOBAE, near *Cuscuta applanata* Engelm. It differs from this species, however, in having longer pedicels and a looser inflorescence and in not exhibiting keels on the calyx. It superficially resembles *Cuscuta umbellata* H. B. K. The obtuse corolla and calyx lobes easily distinguish this species, however.

The only specimen examined was from Durango, Mexico (*Endlich 268*), the type, in the herbaruim of the Botanical Institute at Dahlem (a small bit in the writer's herbarium).

DEPAUW UNIVERSITY

INDEX TO AMERICAN BOTANICAL LITERATURE

1908-1921

The aim of this Index is to include all current botanical literature written by Americans, published in America, or based upon American material; the word America being used in the broadest sense.

Reviews, and papers that relate exclusively to foresty, agriculture, horticulture, manufactured products of vegetable origin, or laboratory methods are not included, and no attempt is made to index the literature of bacteriology. An occasional exception is made in favor of some paper appearing in an American periodical which is devoted wholly to botany. Reprints are not mentioned unless they differ from the original in some important particular. If users of the Index will call the attention of the editor to errors or omissions, their kindness will be appreciated.

This Index is reprinted monthly on cards, and furnished in this form to subscribers at the rate of three cents for each card. Selections of cards are not permitted; each subscriber must take all cards published during the term of his subscription. Correspondence relating to the card issue should be addressed to the Treasurer of the Torrey Botanical Club.

Acosta, C. Colección de maderas cubanas. Revista Agr. Com. y Trab. **3**: 55. F 1920.

Acosta, C. El tararaco [*Hippeastrum*]. Revista Agr. Com. y Trab. **3**: 56. F 1920. [Illust.]

Acosta, C. Sobre el cayeput [*Melaleuca leucadendron* L.] Revista Agr. Com. y Trab. **2**: 535–537. *f. 1, 2.* N. 1919

Addis, J. M. Estudio de los platanos y guineos cubanos. Revista Agr. Com. y Trab. **3**: 418–429. *f. 1–28.* D 1920.

Andrews, A. L. Notes on North American *Sphagnum.* VI. Bryologist **18**: 1–6. 16 Ja 1915.

Andrews, A. L. Notes on North American *Sphagnum.* VII. Bryologist **20**: 84–89. 13 O 1917.

Andrews, A. L. *Tortula caroliniana*, new species. Bryologist **23**: 72–76. *pl. 5.* 2 D 1920.
From North Carolina.

Arber, A. Leaves of certain amaryllids. Bot. Gaz. **72**: 102–105. *f. 1–8.* 15 Au 1921.

Arber, A. The leaf structure of the Iridaceae, considered in relation to the phyllode theory. Ann. Bot. **35**: 301–336. *f. 1–66.* Jl 1921.

Ball, C. R. The relation of crop-plant botany to human welfare. Am. Jour. Bot. **8:** 323–338. Jl 1921.

Ballou, C. H. Los mejores mangos para Cuba. Revista Agr. Com. y Trab. **1:** 187–191. Ap 1918. [Illust.]

Bennett, A. *Sparganium angustifolium* Michx. Jour. Bot. **59:** 235, 236. Au 1921.

Bergman, H. F. Intra-ovarial fruits in *Carica Papaya*. Bot. Gaz. **72:** 97–101. *f. 1–6.* 15 Au 1921.

Berry, E. W. A fossil sea bean from Venezuela. Am. Jour. Sci. **50:** 310–313. *f. 1.* O 1920.

Berry, E. W. Age of the Dakota flora. Am. Jour. Sci. **50:** 387–390. N 1920.

Berry, E. W. A palm nut from the Miocene of the Canal Zone. Proc. U. S. Nat. Mus. **59:** 21, 22. *f. 1–3.* 1921.

Berry. E. W. Paleobotany: a sketch of the origin and evolution of floras. Smithsonian Report **1918:** 289–407. *pl. 1–6* + *f. 1–42.* 1921.

Berry, E. W. Tertiary fossil plants from Costa Rica. Proc. U. S. Nat. Mus. **59:** 169–185. *pl. 22–27.* 1921.
Includes 8 new species.

Blakeslee, A. F., Welch, D. S., & Cartledge, J. L. Technique in contrasting mucors. Bot. Gaz. **72:** 162–172. *f. 1–2.* 15 S 1921.

Blasquez, A. La vegetacion las lluvias. Revista Agricola **6:** 333–345. Je 1920.

Bois, D., & Gerôme, J. La chilacayote du Mexique (courge de Siam): *Cucurbita ficifolia* Bouché (*C. melanosperma* Al. Braun). Bull. Mus. Hist. Natur. **26:** 675–678. 1920.

Borden, A. D. A biological study of the red date-palm scale. *Phoenicococcus Marlatti.* Jour. Agr. Research **21:** 659–667. *pl. 127–130.* 1 Au 1921.

Botto, A. La sweet tussac, *Phalaris bulbosa* Cav. Revista Facult. Agron. y Veter. Univ. La Plata II. **12:** 226–242. *f. 1–4.* 1916; **13:** 1–76. *f. 1–9.* 1918.

Britton, N. L. Stewardson Brown. Jour. N. Y. Bot. Gard. **22:** 110–112. Je 1921.

Bruner, S. C. Enfermedades de la vid [*Vitis*] en Cuba. Revista Agr. Com. y Trab. **1:** 406–409. Au 1918. [Illust.]

Bruner, S. C. La enfermedad del "mosaico" o de "Rayas amarillas" de la caña de azucar en Cuba. Revista Agr. Com. y Trab. **2**: 437–441. *f. 1, 2.* S 1919.

Bruner, S. C. La "*Phomopsis*" de la berenjena. Revista Agr. Com. y Trab. **1**: 468, 469. S 1918.

Bruner, S. C. La pudrición negra del cacao. Revista Agr. Com. y Trab. **2**: 630–636. D 1919. [Illust.]

Calvino, M. El zacate prodigio (*Tripsacum latifolium*, Hitchcock). Revista Agr. Com. y Trab. **3**: 62–67. F 1920. [Illust.]

Calvino, M. Estudio sobre el cultivo de la soya [*Soja max*] en Cuba. Revista Agr. Com. y Trab. **3**: 124–131. My 1920. [Illust.]

Calvino, M. La lechuga [*Lactuca sativa*], su cultivo y aprovechamiento. Revista Agr. Com. y Trab. **1**: 411–418. Au 1918. [Illust.]

Calvino, M. La luz y la producción vegetal. Revista Agr. Com. y Trab. **3**: 532–536. Ap 1921. [Illust.]

Calvino, M. La jicama de agua (*Pachyrhizus tuberosus*) Revista Agr. Com. y Trab. **2**: 84–87. Mr 1919. [Illust.]

Calvino, M. Nuevos variedades de caña de azúcar. Rev. Agr. Com. y Trab. **3**: 436–440. Ja 1921. [Illust.]

Carpenter, C. W. Morphological studies of the *Pythium*-like fungi associated with root rot in Hawaii. Bull. Exp. Sta. Hawaiian Sugar Pl. Assoc. Bot. **3**: 59–65. *pl. 16–23.* Au 1921.

Child, C. M. Le probleme de l'integration physiologique. Scientia **30**: 115–126. 1 Au 1921.

Clute, W. N. Note and comment. Am. Bot. **27**: 116–113. Au 1921.

Clute, W. N. The Victoria water lily. Am. Bot. **27**: 81–86. Au 1921.

Collins, J. L. Reversion in composites. Jour. Hered. **12**: 129–133. *f. 16–19* 10 Au 1921.

Conard, H. S. Fossil plants and classification. Am. Bot. **27**: 95–101. Au 1921.

Coulter, J. M., & Land, W. J. G. A homosporous American *Lepidostrobus*. Bot. Gaz. **72**: 106–108. 15 Au 1921.

Cremata, M. Una excursión botanica a Isla de Pinos. Revista Agr. com. y Trab. **3**: 47–49. F 1920. [Illust.]

Cruz, F. B. El cultivo del tabaco en Cuba. Revista Agr. Com. y Trab. **1**: 172–183. Ap 1918. [Illust.]

Cummings, C. E. Mushrooms. Hobbies **2**: 3–17. S 1921.

Dachnowski, A. P. Peat deposits and their evidence of climatic changes. Bot. Gaz. **72**: 57–89. *f. 1–12.* 15 Au 1921.

Davidson, A. New species from southern California. Bull. S. California Acad. Sci. **20**: 49–53. Au 1921 [Illust.]

Dusen, P., & Neger; F. W. Über Xylopodien. Beih. Bot. Centralbl. **38**: 258–317. *f. 1–10.* 25 My 1921.
Studies of Brazilian plants.

Duursma, G. D. Pilocereussen. Floralia **42**: 376–378. 17 Je 1921. [Illust.]

Duursma, G. D. Rhipsalideae. Floralia **42**: 378, 379. 17 Je 1921.

Eames, E. A. An unusual form of *Habenaria clavellata.* Rhodora **23**: 126–127. *pl. 131.* 26 Au 1921.

Edgerton, C. W., & Tiebout, G. L. The mosaic disease of the Irish potato and the use of certified potato seed. Louisiana Bull. 181: 1–15. *f. 1–3.* Au 1921.

Elorduy, S. T. Selección del tabaco. Rev. Agricola **5**: 801–810. *f. 1–9.* Ap 1921; **6**: 14–18. *f. 10–12.* My 1921.

Emerson, R. A The genetic relations of plant colors in maize. Cornell Univ. Agr. Exp. Sta. Mem. 39: 1–156. *pl. 1–11.* Mr 1921.

Felippone, F. Contribution à la flore bryologique de l'Uruguay. Fasc. 3: 40–45. 1917. [Illust.]
Includes new species in *Didymodon* (1), *Leptodontium* (1), *Fabronia* (1).

Feustel, H. Anatomie und Biologie der Gymnospermenblätter. Beih. Bot. Centralbl. **38**: 177–257. 25 My 1921.

Fortun, G. M. Notas sobre una excursion a "El Retiro" [Cuba]. Revista Agr. Com. y Trab. **3**: 410–413. D 1920. [Illust.]
Residence of Jose Blain.

Fortun, G. M., & Bruner, S. C. Investigaciones sobre la enfermedad del "mosaico" o "rayas amarillas" dela caña de azucar. Revista Agr. Com. y Trab. **3**: 441–445. Ja 1921. [Illust.]

Fraser, G. The dwarf trees of Vancouver Island. Gard. Chron. **70**: 102. *f. 40.* 20 Au 1921.

Fred, E. B. The fixation of atmospheric nitrogen by inoculated soybeans. Soil. Sci. **11**: 469–472. *pl. 1–3.* Je 1921.

Fred, E. B., Wright, W. H., & Frazier, W. C. Field tests on the inoculation of canning peas. Soil Sci. **11**: 479–491. *pl. 1–3.* Je 1921.

Frye, T. C. Notes on useful and harmful mosses. Bryologist **23**: 71. 2 D 1920.

Gagnepain, F. Classification des *Eugenia.* Bull. Soc. Bot. France **64**: 94–103. 1917.

Gile, P. L., & Carrero, J. O. Assimilation of nitrogen, phosphorus and potassium by corn when nutrient salts are confined to different roots. Jour. Agr. Research **21**: 545–573. *f. 1.* 15 Jl 1921.

Girola, C. D. Contribución al curso de cultivas industriales. Revista Facult. Agron. y Veter. Univ. La Plata II. **10**: 83–113. 1913.

Girola, C. D. El cultivo de la yerba-mate. (*Ilex paraguariensis* St. Hil.) Revista Facult. Agron. y Veter. Univ. La Plata II. **11**: 163–184. *f. 1–4.* 1915; **12**: 196–225. *f. 1–6.* 1916.

Girola, C. D. Monografia sobre el cañamo (*Cannabis sativa* L.). Revista Facult. Agron. y Veter. Univ. La Plata II. **10**: 9–68. *f. 1–13.* 1913.

Girola, C. D. Observaciones sobre el cultivo del henequen. Revista Facult. Agron. y Veter. Univ. La Plata II. **11**: 101–151. *f. 1–19.* 1915.

Grant, M. Saving the redwoods. Nat. Geog. Mag. **37**: 519–536. Je 1920.

Griffiths, D. The production of the Easter lily in northern climates. U. S. Dept. Agr. Bull. 962: 1–31. *f. 1–14.* 12 Au 1921.

Grosovich, J. M. Las principales malezas de los sembrados. Revista Zootech. **8**: 186–192. 15 My 1921.

Grove, W. B. Species placed by Saccardo in the genus *Phoma.* Kew Bull. Misc. Inform. **1921**: 136–157. 1921.
Includes 2 new American combinations in *Phomopsis.*

Hardy, M. E. Earth stars. Am. Bot. **27**: 86, 87 Au 1921.

Harris, J. A. Leaf-tissue production and water content in a mutant race of *Phaseolus vulgaris.* Bot. Gaz. **72**: 151–161. 15 S 1921.

Harter, L. L., & Weimer, J. L. Studies in the physiology of parasitism with special reference to the secretion of pectinase by *Rhizopus Tritici*. Jour. Agr. Research **21**: 609–625. 1 Au 1921.

Heim, A. Charakterpflanzen der Halbinsel Niederkalifornien. Vegetationsbilder XIII. **3, 4**: *pl. 13–24* + descriptive text. 1916.

Hoerner, G. R. Germination of aeciospores, urediniospores and teliospores of *Puccinia coronata*. Bot. Gaz. **72**: 173–177. 15 S 1921.

Hohenkerk, L. S. Botanical identifications of British Guiana trees and plants. Jour. Board Agr. British Guiana **11**: 98–106, 178–185. 1918.

Hohenkerk, L. S. British Guiana timbers. Jour. Board Agr. British Guiana **12**: 152–187. Jl 1919.

Hollick, A. Loco weeds. Nat. Hist. **21**: 85–91. F 1921.

H[osseus], C. C. Dr. Federico Kurtz [1854–1920]. Bol. Acad. Nac. Cienc. Cordoba (Argentina) **24**: LIV–LIX. 1921.

Howe, M. A. Some plants from tropical sea gardens. Nat. Hist. **20**: 561–568. D 1920.

Hutchinson, J. The family Winteraceae. Kew Bull. Misc. Inform. **1921**: 185–191. 1921. [With map.]

Hutchinson, J. The genus *Therorhodion*. Kew Bull. Misc. Inform. **1921**: 201–205. *f. 1*. 1921.

Illick, J. S. The hard pines of the northeast. Am. For. **27**: 487–496. Au 1921.

Illick, J. S. The pines of the South. Am. For. **27**: 551–559, 574. S 1921. [Illust.]

Johnston, J. R. El platano y sus enfermedades. Revista Agr. Com. y Trab. **1**: 419–421. Au 1918. [Illust.]

Jones, L. H., & Shive, J. W. Effect of ammonium sulphate upon plants in nutrient solutions supplied with ferric phosphate and ferrous sulphate as sources of iron. Jour. Agr. Research **21**: 701–728. *pl. 136* + *f. 1–9*. 15 Au 1921.

Jones, S. R. Preliminary report on the flora and fauna of the eastern part of the Rosebud Reservation, now known as Gregory County [South Dakota]. South Dakota Geol. Surv. Bull. **4**: 123–142. 1908.

Jouvenaz, J. H. *Cereus*. Floralia **42**: 371, 372. 17 Je 1921. [Illust.]

Jouvenaz, J. H. *Echinocactus.* Floralia **42**: 372. 17 Je 1921. [Illust.]

Jouvenaz, J. H. *Echinocereus.* Floralia **42**: 372. 17 Je 1921. [Illust.]

Kearney, T. H. Pollination of Pima cotton in relation to the yield of seed and fiber. Jour. Hered. **12**: 99–101. 10 Au 1921.

Kermode, F. A preliminary catalogue of the flora of Vancouver and Queen Charlotte Islands. Pp. 1–89. Victoria, British Columbia. 1921.

Kirjasoff, A. B. Formosa, the beautiful. Nat. Geog. Mag. **37**: 247–292. Mr 1920.

Kobel, F. Einige Bemerkungen zu den *Astragalus*- und *Cytisus*-bewohnenden *Uromyces*-Arten. Ann. Mycol. **19**: 1–16. 1921.

Korstian, C. F. Grazing practice on the natural forests and its effect on natural conditions. Sci. Mo. **13**: 275–281. *f. 1–7.* S 1921.

Kräusel, R. Die Bedeutung der Anatomie lebender und fossiler Hölzer für die Phylogenie der Koniferen. Naturw. Wochenschr. **32**: 305–311. *f. 1–9.* 10 Je 1917.

Kräusel, F. Welche Ergebnisse liefert die Untersuchung tertiärer Pflanzenreste? Naturw. Wochenschr. **33**: 209–213. 14 Ap1918.

Krieger, L. C. C. Common mushrooms of the United States. Nat. Geog. Mag. **37**: 387–439. *pl. 1–16* My 1920.

Lantes, A. El alamo [*Populus*]. Revista Agr. Com. y Trab. **2**: 612, 613. D 1919. [Illust.]

Lingelsheim, A. Oleaceae-Oleoideae-Fraxineae. Pflanzenreich IV. 243: 1–66, 115–125. *f. 1–11* + *map.* 29 Je 1920.
Includes 2 new North American species of *Fraxinus.*

Livingston, B. E., & Shreve, F. The distribution of vegetation in the United States, as related to climatic conditions. Carnegie Inst. Washington Publ. 284: I–XVI + 1–590. *pl. 1–73* + *f. 1–74.* 1921.

Locy, W. A. The earliest printed illustrations of natural history. Sci. Mo. **13**: 238–258. *f. 1–10.* S 1921.

MacDougal, D. T. Growth in trees. Carnegie Inst. Washington Publ. 307: 3–41. *f. 1–16.* 1921.

MacDougal, D. T. Growth in trees. Proc. Am. Philos. Soc. **60:** 7–14. *pl. 1 + f. 3–5.* 1921.

MacDougal, D. T. The action of bases and salts on biocolloids and cell-masses. Proc. Am. Philos. Soc. **60:** 15–30. 1921.

MacDougal, D. T., & Working, E. B. Another high-temperature record for growth and endurance. Science II. **54:** 152, 153. 19 Au 1921.

McMurray, N. Freak dogwood. Am. Bot. **27:** 113. Au 1921.

MacTaggart, A. The influence of certain fertilizer salts on the growth and nitrogen-content of some legumes. Soil Sci. **11:** 435, 454. *pl. 1.* Je 1921.

Meier, H. F. A. Effect of direct current on cells of root tip of Canada field pea. Bot. Gaz. **72:** 113–138. *pl. 2–3 + f. 1–3.* 15 S 1921.

Moodie, R. L. Bacteria in the American Permian. Science II. **54:** 194–196. 2 S 1921.

Mottier, D. M. On certain plastids, with special reference to the protein bodies of *Zea*, *Ricinus*, and *Conopholis*. Ann. Bot. **35:** 349–364. *pl. 15.* Jl 1921.

[**Nicholls, H. A. A.**] Useful timbers of Dominica. In Bell, H. H., Notes on Dominica and hints to intending settlers. 34–56. 1919. [Illust.]

Nichols, G. E. The vegetation of Connecticut. VII. The associations of depositing areas along the seacoast. Bull. Torrey Club **47:** 511–548. *f. 1–10.* 24 N 1920.

Oakley, R. A. Effect of the length of day on seedlings of alfalfa varieties and the possibilities of utilizing this as a practical means of identification. Jour. Agr. Research **21:** 599–607. *pl. 111–121.* 15 Jl 1921.

Overton, J. B. The mechanism of root pressure and its relation to sap flow. Am. Jour. Bot. **8:** 369–374. Jl 1921.

Pack, D. A. Chemistry of after-ripening, germination and seedling development of juniper seeds. Bot. Gaz. **72:** 139–150. 15 S 1921.

Pellett, F. C. American honey plants. pps. 1–297. *f. 1–152.* 1920.

Pennington, L. H., & others. Investigations of *Cronartium ribicola* in 1920. Phytopathology **11:** 170–172. Ap 1921.

Petrak, F. Mykologische Notizen. II. Ann. Mycol. **19:** 17–128. 1921.

Pittier, H. Acerca del genero *Gyranthera* Pittier. Contrib. Fl. Venezuela 14–19. 1921.
Includes *G. caribensis* sp. nov. from Venezuela.

Pittier, H. Arbdes y arbustos nuevos de Venezuela. I. Contrib. Fl. Venezuela 3–14. 1921.
New species in *Inga* (1), *Raputia* (1), *Esenbeckia* (1), *Trichilia* (1), *Stillingia* (1), *Gustavia* (1), *Mouriria* (1), *Oxythece* (1), *Pouteria* (1), *Sideroxylum* (1).

Plemper van Balen, B. A. *Cephalocereus senilis*, de "Grijsaard." Floralia **42:** 371. 17 Je 1921. [Illust.]

Pondal, M. L. Fabricaciòn del extracto de quebracho. Bol. Acad. Nac. Cienc. Cordoba (Argentina) **24:** 293–324. *f. 1–14.* 1921.

Popenoe, W. La mora gigante de Colombia. Revista Agricola **6:** 503–507. S 1920.

Popenoe, W. The pejibaye, a neglected food-plant of tropical America. Jour. Hered. **12:** 154–166 *f. 4–9.* Ap 1921.

Povah, A. H. W. An attack of poplar canker following fire injury. Phytopathology **11:** 157–165. *f. 1–3.* Ap 1921.

Prouty, W. F. A more phenomenal shoot. Science II **54:** 170. 26 Au 1921.
Shoot of *Paulownia tomentosa.*

Ramírez, R. El chahiuztle rojo del frijol. Revista Agricola **5:** 830. Ap 1921. [Illust.]

Ramírez, R. La cenicilla del tomate, *Physalis dubescens* L. Revista Agricola **5:** 830. Ap 1921. [Illust.]

Reagan, A. B. Notes on the flora and fauna of the Rosebud Reservation west of Gregory County [South Dakota]. South Dakota Geol. Surv. Bull. **4:** 143–167. 1908.

Reiche, K. Die physiologische Bedeutung des anatomischen Baues der Crassulaceen. Flora **114:** 249–255. *f. 1, 2.* 1921.

Reiche, K. Zur Kenntnis von *Sechium edule* Sev. Flora **114:** 232–248. *f. 1–9.* 1921.

Reiche, K. Zur Kenntnis von *Senecio praecox* DC. Flora **114:** 255–261. *f. 3, 4.* 1921.

Renson, C. El *Tripsacum latifolium*, Hitchc. Revista Agricola **5:** 351–368. Je 1919.

Richards, V. F. The sassafras. Am. Bot. **27:** 105. Au 1921.

Roberts, H. F. Relation of hardness and other factors to protein content of wheat. Jour. Agr. Research **21**: 507–522. *pl. 100.* 15 Jl 1921.

Rock, J. F. The Okala berry of Hawaii. Jour. Hered. **12**: 147–150. *f. 1, 2.* Ap 1921.

Roig, J. F., Cremata, M., & Bruner, S. C. Exploración botanica en la Cienaga de Zapata [Cuba]. Revista Agr. Com. & Trab. **3**: 213–221. Jl. 1920. [With map.]

Safford, W. E. *Datura*, an inviting genus for the study of heredity. Jour. Hered. **12**: 178–190. *f. 10–16.* Ap 1921.

Shufeldt, R. W. Common American mushrooms. Am. For. **27**: 579–587. *f. 1–13.* S 1921.

Smith, E. F. Effect of crowngall inoculations on *Bryophyllum.* Jour. Agr. Research **21**: 593–597. *pl. 101–110.* 15 Jl 1921.

Snell, W. H. Chlamydospores of *Fomes officinalis* in nature. Phytopathology **11**: 173–175. *f. 1.* Ap 1921.

Sprague, T. A. A revision of the genus *Capraria.* Kew Bull. Misc. Inform. **1921**: 205–212. 1921.

Sprague, [T. A.]. *Begonia (Gireoudia) trigonoptera*, Sprague [Begoniaceae]. Kew Bull. Misc. Inform. **1921**: 218, 219. 1921.
New species from Guatemala.

S[prague], T. A. The generic name *Wikstroemia.* Kew Bull. Misc. Inform. **1921**: 175, 176. 1921.

Sprague, T. A., & Riley, L. A. M. Notes on *Raimannia* and allied genera. Kew Bull. Misc. Inform. **1921**: 198–201. 1921.
Includes nine new combinations.

Taylor, N. The flora above the timber line of Mount Marcy. Am. Scien. & Hist. Preserv. Soc. Ann. Rep. **25**: 317, 318. 1920.

Thatcher, L. E. A fungus disease suppressing expression of awns in a wheat-spelt hybrid. Jour. Agr. Research **21**: 699, 700. *pl. 135.* 15 Au 1921.

Theye, C. Nutrición nitrogenada de los vegetales. Revista Facult. Letr. y Cienc. Univ. Habana **31**: 145–151. Je 1921.

Thurston, H. W., & Orton, C. R. A *Phytophthora* parasitic on peony. Science II. **54**: 170, 171. 26 Au 1921.

Tisdale, W. H. Two *Sclerotium* diseases of rice. Jour. Agr. Research **21**: 649–657. *pl. 122–126.* 1 Au 1921.

Torres, L. G. Plantas forrajeras: una leguminosa forrajera indigena (*Desmodium leiocarpum* G. Don—*Meibomia leiocarpa*). Revista Agricola **6**: 84–86. Je 1921. [Illust.]

Trelease, W. A natural group of unusual black oaks. Proc. Am. Philos. Soc. **60**: 31–33. *pl. 2–4.* 1921.
Includes 3 new species from Mexico.

Uphof, J. C. T. Vegetationsbilder aus dem Staate Michigan. Vegetationsbilder **13**: *pl. 43–48.* 1921.

Valdivia Montanez, M. A. El mani, su cultivo, productos y usos. Revista Agr. Com. y Trab. **1**: 570–579. N 1918. [Illust.]
Arachis hypogea L.

Valencia, F. V. Mechanical tests of some commercial Philippine timbers. Philipp. Jour. Sci. **18**: 485–535. *pl. 1 + f. 1–19.* My 1921.

Vendrell, E. La selección de las semillas. Revista Agr. Com. y Trab. **1**: 260, 261. My 1918.

Verbeek Wolthuys, J. J. Pereskieae en Opuntieae. Floralia **42**: 373–375, 387–389. Je 1921. [Illust.]

Visher, S. S. The geography of South Dakota. South Dakota Geol. & Nat. Hist. Surv. Bull. **8**: 1–189. *f. 1–52.* Jl 1918.
Contains chapter on biogeography. [Corrected reprint.]

Waby, J. Z. Some interesting species of palms. Jour. Board Agr. British Guiana **11**: 172, 173. O 1918; **12**: 188–195. Jl 1919.

Weimer, J. L., & Harter, L. L. Respiration and carbohydrate changes produced in sweet potatoes by *Rhizopus Tritici*. Jour. Agr. Research **21**: 627–635. 1 Au 1921.

Weimer, J. L., & Harter, L. L. Wound-cork formation in the sweet potato. Jour. Agr. Research **21**: 637–647. 1 Au 1921.

Weir, J. R. Note on *Cenangium abietis* (Pers.) Rehm on *Pinus ponderosa*. Phytopathology **11**: 166–170. *f. 1.* Ap 1921.

Weir, J. R., & Hubert, E. E. Forest disease surveys. U. S. Dept. Agr. Bull. 658: 1–23. *f. 1–23.* 12 Je 1918.

Weiss, F., & Harvey, R. B. Catalose hydrogen-ion concentration and growth in the potato wart disease. Jour. Agr. Research **21**: 589–592. 15 Jl 1921.

Welton, F. A. Longevity of seeds. Monthly Bull. Ohio Agr. Exp. Sta. **6**: 18–24. 1921. [Illust.]
[Corrected reprint.]

Wercklè, C. Die natürlichen Wachstumsbedingungen der epiphytischen Orchideen in Costa Rica. Gartenflora **70**: 90–94. 15 My 1921.

Wester, J. P. The mango. Philipp. Bur. Agr. Bull. **18**: 1–70. *pl. 1–17*. 1920.
Second revised edition.

Wherry, E. T. The reactions of the soils supporting the growth of certain native orchids. Jour. Washington Acad. Sci. **8**: 591–598. 4 N 1918.

Wherry, E. T. Soil reactions of *Spiranthes cernua* and its relatives. Rhodora **23**: 127–129. 26 An 1921.

White, D. Value of floral evidence in marine strata as indicative of nearness of shores. Bull. Geol. Soc. Amer. **22**: 221–227. 10 My 1911.

Williams, R. S. *Hyophila subcucullata* sp. nov. Bryologist **24**: 22. *pl. 2*. 25 Jl 1921.
From Cuba.

Williams, R. S. *Sematophyllum Smallii* sp. nov. Bryologist **23**: 76–78. *pl. 6*. 2 D 1920.
From Florida.

Wilson, W. J. Palaeobotany. Summ. Rep. Canada Geol. Surv. Dept. Mines **1916**: 300–302. 1917.

Wright, C. H. *Aechmea chromatica*, C. H. Wright [Bromeliaceae-Bromelieae]. Kew Bull. Misc. Inform. **1921**: 220, 221. 1921.
New species from tropical America.

Yamanouchi, S. Life history of *Corallina officinalis* var. *mediterranea*. Bot. Gaz. **72**: 90–96. 15 Au 1921.